

Spitzer POWER-LAW AGN CANDIDATES IN THE *Chandra* DEEP FIELD-NORTH

J. L. DONLEY,¹ G. H. RIEKE,¹ P. G. PÉREZ-GONZÁLEZ,^{1,2} J. R. RIGBY,¹ A. ALONSO-HERRERO³

Accepted for publication in *The Astrophysical Journal*

ABSTRACT

We define a sample of 62 galaxies in the *Chandra* Deep Field-North whose *Spitzer* IRAC SEDs exhibit the characteristic power-law emission expected of luminous AGN. We study the multiwavelength properties of this sample, and compare the AGN selected in this way to those selected via other *Spitzer* color-color criteria. Only 55% of the power-law galaxies are detected in the X-ray catalog at exposures of > 0.5 Ms, although a search for faint emission results in the detection of 85% of the power-law galaxies at the $\geq 2.5\sigma$ detection level. Most of the remaining galaxies are likely to host AGN that are heavily obscured in the X-ray. Because the power-law selection requires the AGN to be energetically dominant in the near- and mid-infrared, the power-law galaxies comprise a significant fraction of the *Spitzer*-detected AGN population at high luminosities and redshifts. The high $24\ \mu\text{m}$ detection fraction also points to a luminous population. The power-law galaxies comprise a subset of color-selected AGN candidates. A comparison with various mid-infrared color selection criteria demonstrates that while the color-selected samples contain a larger fraction of the X-ray luminous AGN, there is evidence that these selection techniques also suffer from a higher degree of contamination by star-forming galaxies in the deepest exposures. Considering only those power-law galaxies detected in the X-ray catalog, we derive an obscured fraction of 68% (2:1). Including all of the power-law galaxies suggests an obscured fraction of $< 81\%$ (4:1).

Subject headings: galaxies: active — infrared: galaxies — X-rays: galaxies

1. INTRODUCTION

Detecting complete samples of AGN, both locally and in the deep cosmological fields, has been a major ongoing goal. Hard X-ray selection is a powerful way to detect both relatively uncontaminated and complete samples of AGN. Deep X-ray surveys with *Chandra* and *XMM-Newton* have now resolved 70-90% of the cosmic X-ray background (CXRB) at 2-8 keV into discreet sources (Mushotzky et al. 2000; Giacconi et al. 2002; Alexander et al. 2003; Bauer et al. 2004; Worsley et al. 2004, 2005), detecting the majority of the X-ray unobscured AGN population in the deep X-ray fields. At high column densities, however, the dust and gas surrounding the central engine (in combination with that located in the host galaxy) are capable of hiding virtually all accessible AGN tracers. Therefore, while the overall resolved fraction of the CXRB is high, it drops with increasing energy to 60% at 6-8 keV and to 50% at > 8 keV (Worsley et al. 2004, 2005).

Population synthesis models of the CXRB and of X-ray luminosity functions therefore predict a significant population of heavily obscured AGN not detected in the deepest X-ray fields (i.e., the 1 Ms *Chandra* Deep Field-South (CDF-S) and the 2 Ms *Chandra* Deep Field-North (CDF-N)). Predictions of their properties are in rough, but not complete, agreement. Treister et al. (2004) predict an X-ray incompleteness of 25% at $N_{\text{H}} = 10^{23}\ \text{cm}^{-2}$ and of 70% at $N_{\text{H}} = 10^{24}\ \text{cm}^{-2}$. Ballantyne et al. (2006) estimate that the deep X-ray surveys miss $\sim 50\%$ of obscured AGN with $\log L_{\text{x}}(\text{ergs s}^{-1}) > 44$ at all z , and that most of the missing objects are Compton-thick ($\log N_{\text{H}}(\text{cm}^{-2}) > 24$). Worsley et al. (2005) find that the unresolved sources are likely to lie at redshifts of $z = 0.5 - 1.5$,

have column densities in excess of $10^{23}\ \text{cm}^{-2}$, and have intrinsic X-ray luminosities of $< 5 \times 10^{43}\ \text{ergs s}^{-1}$.

Numerous attempts have been made to detect this population of heavily obscured AGN, many of which have focused on the mid-infrared (MIR) emission where the obscured radiation is re-emitted (Ivison et al. 2004; Lacy et al. 2004; Hatziminaoglou et al. 2005; Stern et al. 2005; Alonso-Herrero et al. 2006; Polletta et al. 2006) or on combinations of MIR and multi-wavelength data (Johansson et al. 2004; Yan et al. 2004, 2005; Donley et al. 2005; Franceschini et al. 2005; Houck et al. 2005; Leipski et al. 2005; Martínez-Sansigre et al. 2005, 2006; Richards et al. 2006; Weedman et al. 2006). In the MIR, luminous AGN can often be distinguished by their characteristic power-law emission, which extends from the infrared to the ultraviolet (e.g. Neugebauer 1979; Elvis et al. 1994). This emission is not necessarily due to a single source, but can arise from the combination of non-thermal nuclear emission and thermal emission from various nuclear dust components (e.g. Rieke & Lebofsky 1981). In contrast, star-forming galaxies are characterized by dust and stellar emission features redward and blueward of a local minimum at $\sim 5\ \mu\text{m}$.

We focus here on AGN with red power-law SEDs in the *Spitzer* 3.6-8.0 μm bands. Alonso-Herrero et al. (2006) selected a sample of 92 such sources in the CDF-S, 70% of which are hyper-luminous infrared galaxies (Hyper-LIRGS, $\log L_{\text{IR}}(L_{\odot}) > 13$) or ultra-luminous infrared galaxies (ULIRGS, $\log L_{\text{IR}}(L_{\odot}) > 12$). Nearly half (47%) of their power-law sample were not detected in X-rays at exposures of up to 1 Ms. We use a selection similar to that of Alonso-Herrero et al. (2006) to identify power-law galaxies in the 2 Ms CDF-N. Because the central regions of the *Chandra* Deep Fields are photon-limited, not background-limited, the 2 Ms CDF-N is twice as sensitive as the 1 Ms CDF-S at the *Chandra* aimpoint (e.g. Alexander et al. 2003).

We use a combination of spectroscopic and photometric redshifts to estimate distances and luminosities for this class

¹ Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721; jdonley@as.arizona.edu

² Departamento de Astrofísica y CC. de la Atmósfera, Facultad de CC. Físicas, Universidad Complutense de Madrid, 28040 Madrid, Spain

³ Departamento de Astrofísica Molecular e Infrarroja, Instituto de Estructura de la Materia, CSIC, E-28006 Madrid, Spain

of AGN, placing them in the context of the overall AGN population. The observations and data reduction are outlined in §2 and the selection criteria are discussed in §3. The photometric redshift code is discussed in §4. Making use of the deepest available (2 Ms) X-ray data, we search for faint X-ray emission from sources missed in the X-ray catalogs. The X-ray, MIR, radio, and optical properties of the sample, including detection fractions, X-ray luminosities, radio classifications, and optical–MIR SEDs, are discussed in §5. In §6, we compare the power-law selection to other MIR color selection techniques, and discuss the completeness and reliability of the power-law and color-selection criteria. Finally, in §7, we use the X-ray data to calculate the obscuring columns of the power-law galaxies and to estimate the obscured fraction of the power-law sample. Throughout the paper, we assume a cosmology of $(\Omega_m, \Omega_\Lambda, H_0) = (0.3, 0.7, 72 \text{ km s}^{-1} \text{ Mpc}^{-1})$.

2. OBSERVATIONS AND DATA REDUCTION

Spitzer was used to obtain MIPS and IRAC MIR observations of the CDF-N, with 1400 s and 500 s exposures, respectively. We use these MIPS GTO images in place of the GOODS *Spitzer* data because the former cover the full CDF-N, whereas the latter cover a field approximately two times smaller. The MIPS image was processed using the MIPS GTO data analysis tool (Gordon et al. 2004, 2005); the IRAC data analysis is described in Huang et al. (2004). We used the IRAF task *allstar* to select sources at 24 μm , and SEXTRACTOR (Bertin & Arnouts 1996) to select sources in the IRAC bands. The 80% completeness limit for the 24 μm data is 83 μJy (Papovich et al. 2004). The IRAC detection limits of our survey (see §3) are 1.8, 2.8, 14.5, and 18.0 μJy , at 3.6, 4.5, 5.8, and 8.0 μm , respectively. Optical and NIR photometry were measured from the GOODS dataset (*bviz*, Giavalisco et al. 2004), as well as from the data of Capak et al. (2004) (*UBVRiz'HK'*). The ground-based photometry was aperture-matched for consistency, using the aperture of the most sensitive band in which the source was detected. For cases in which multiple ground-based sources in a 1.5'' radius caused the aperture-matched flux density to differ from that of the nearest source by a factor of ≥ 1.5 , we replaced the aperture-matched flux densities with the cataloged flux densities of the nearest source in all bands. In addition, if the ground-based photometry of sources with multiple GOODS counterparts (in a 1.5'' radius) differed from the GOODS photometry by a factor of 2 or more in any of the bands, we treated the ground-based photometry as upper limits. For further details, see Pérez-González et al. (2005).

3. SAMPLE SELECTION

We selected as power-law galaxies sources that were detected with $S/N \geq 6$ in each of the four IRAC bands and whose IRAC spectra are well-fit by a line of spectral index $\alpha \leq -0.5$, where $f_\nu \propto \nu^\alpha$. In a similar study, Alonso-Herrero et al. (2006) chose a limit of $\alpha \leq -0.5$ based on the mean spectral slope of optically-selected quasars ($\alpha \sim -1$, Elvis et al. 1994; Neugebauer et al. 1979) and the optical spectral indices of Sloan Digital Sky Survey (SDSS) QSOs ($\alpha = 0.5$ to -2 , Ivezić et al. 2002). For comparison, broad-line AGN (BLAGN) from the AGN and Galaxy Evolution Survey (AGES, C. S. Kochanek et al., in prep) have 3.6–8 μm slopes of $\alpha = -1.07 \pm 0.53$ (Stern et al. 2005), indicating that our cut lies within $\sim 1\sigma$ of the mean. To ensure a good linear fit, we applied a cut in the chi-squared probability of $P_\chi \geq 0.1$. P_χ is the probability that a fit to a power-law distribution would

yield a value greater than or equal to the observed chi-squared; a probability of 0.5 corresponds to a reduced chi-squared of 1. P_χ tends to either lie close to 0.5, or is very small (see Bevington & Robinson 2003). Because we are primarily interested in the X-ray and radio properties of the power-law galaxies, we also restricted the sample to those galaxies with X-ray exposures of ≥ 0.5 Ms (Alexander et al. 2003); this selection also ensures deep 1.4 GHz radio coverage (Richards 2000) and results in a total survey area of 350 arcmin². In addition, we required the IRAC SED to rise monotonically in f_ν to prevent contamination from star-forming galaxies with possible stellar features in the IRAC bands (resulting in the rejection of 10 sources). Using these criteria, we identified 79 power-law galaxies in the CDF-N.

We removed 10 of the selected galaxies due to blended or problematic IRAC photometry and we removed 1 galaxy (CDFN 22363) because of its stellar-dominated spectrum. CDFN 22363 has a shallow slope of only -0.56 ± 0.20 and hence might have been scattered into the power-law sample by noise (it has the second flattest slope in the sample). It is the only source in the power-law sample with a spectroscopic redshift that is not detected in X-rays, and it lacks a 24 μm counterpart (see §5.2). Six additional sources were removed because their optical–MIR SEDs exhibited possible stellar bumps. The final sample of 62 power-law galaxies is listed in Table 1; radio through X-ray SEDs are shown in Figure 1. We indicate in Figure 1 the 27 sources that lie in the GOODS ACS region. In addition, we flag all sources for which we have either replaced the ground-based aperture-matched flux densities with the cataloged flux densities of the nearest source or for which we treat the ground-based photometry as upper limits (see §2).

The S/N cut of 6 in each of the IRAC bands imposes the following detection limits on the IRAC 3.6, 4.5, 5.8, and 8.0 μm bands: 1.8, 2.8, 14.5, and 18.0 μJy . This large range of flux limits imposes complicated selection effects on the power-law sample. At $\alpha < -0.67$, the sample is flux-limited by the 5.8 μm band. That is, all sources whose 5.8 μm flux density exceeds the detection limit of 14.5 μJy are detectable in all of the IRAC channels. At $\alpha > -0.67$, however, the limiting flux shifts to the 8.0 μm band. Because there is little change in the 5.8 μm detection limit for red power-law sources with $-0.67 < \alpha < -0.5$ (f_{lim} increases from 14.5 μJy at $-3 < \alpha < -0.67$ to 15.3 μJy at $\alpha = -0.5$), the power-law sample is essentially flux-limited in the 5.8 μm band. Comparisons between the number of red ($\alpha \leq -0.5$) and blue ($\alpha > -0.5$) sources in the full IRAC sample, however, will be complicated by these effects, as the 5.8 μm limiting flux increases with increasing spectral slope, reducing the number of blue sources that meet the requirements of the power-law sample.

To minimize the chances of selecting non-active galaxies via the power-law selection, we compared the spiral galaxy templates from Devriendt et al. (1999) to the optical–MIR SEDs of the sources. None of the sources in the final sample are well-fit by a spiral template. In addition to contamination from local spiral galaxies, Alonso-Herrero et al. (2006) found that cool ULIRGS at $z > 2$ and with $\alpha > -1$ could appear as power-law galaxies. Of the 62 power-law galaxies in the sample, nine have $\alpha > -1$ and potentially lie at $z > 2$. Seven of these sources, however, have X-ray counterparts, and all but one (a source with no redshift estimate) have X-ray luminosities of $\log L_x (\text{ergs s}^{-1}) > 43$, and are therefore unlikely to be starburst-powered ULIRGS. We further explore the possibil-




FIG. 1.— X-ray through radio observed-frame SEDs of the power-law galaxies. The flags printed under the source ID indicate by x whether (1) the source lies in the GOODS ACS field, (2) the aperture-matched flux densities have been replaced by the cataloged flux densities of the nearest source and (3) the ground-based photometry have been treated as upper limits (see §2).

ity of contamination due to star-forming galaxies in §6.2.2, using several updated SEDs, described in detail in Appendix A.

We have also considered whether the contributions of 3.3 μm ($z > 0.5$) and 6.2 μm ($z < 0.5$) aromatic emission features to the flux in the IRAC 8.0 μm band could create the appearance of a red power-law continuum. Observations of starbursts detect these aromatic features to equivalent widths of $\sim 0.1 \mu\text{m}$ and $\sim 0.7 \mu\text{m}$, respectively (Risaliti et al. 2006; B. R. Brandl et al., in preparation). Due to the large bandpass of the 8.0 μm IRAC channel, however, the 3.3 and 6.2 μm emission features should have at most a 3% or 20% effect, respectively, on the total flux. (The models of Devriendt et al. (1999) predict a slightly larger (20%) effect due to the 3.3 μm feature (Stern et al. 2005)). Given the small impact of these features and the small number of power-law galaxies

with $z < 0.5$ for which the 8 μm feature could have an effect, we expect little contamination in the sample.

As a final check on the selection criteria, we plot in Figure 2 the ratio of X-ray to optical emission, a well-known AGN diagnostic (e.g. Maccacaro et al. 1988; Barger et al. 2003; Hornschemeier et al. 2003). Sources not detected in R-band are assigned a lower limit of 26 mag, the approximate completeness limit of the R-band catalog of Capak et al. (2004). As was found by Alonso-Herrero et al. (2006), all of the power-law galaxies lie within the region typically populated by AGN or transition objects, given the current X-ray and optical limits. Four of the X-ray sources appear to be optically-faint with respect to their X-ray fluxes (e.g. Rigby et al. 2005).




FIG. 1.— Continued.

4. REDSHIFTS

Twenty one of the power-law galaxies have secure spectroscopic redshifts (Barger et al. 2001, 2002, 2003; Hornschemeier et al. 2001; Dawson et al. 2003; Cowie et al. 2004; Swinbank et al. 2004; Wirth et al. 2004 [Team Keck Redshift Survey]; Chapman et al. 2005; the Sloan Digital Sky Survey [SDSS]) and 2 additional sources have photometric redshifts from Barger et al. (2003). The spectroscopic redshifts range from $z = 0.29$ to 2.92 , with a mean of $z = 1.77$ and a median of $z = 2.02$. Of these 21 sources, 15 are classified as broad-line AGN, 4 have low S/N narrow emission lines, and 2 are AGN whose line width is not classified in the available literature.

We supplement the available spectroscopic redshifts with photometric redshifts. Photometric redshifts were estimated with an improved version of the method described in Pérez-González et al. (2005). Our technique is based on the construction of a complete set of SED templates composed

from the galaxies with highly reliable spectroscopic redshifts (about 1500 sources selected in the CDF-N and CDF-S), which are later used to fit the SEDs of all the sample and estimate a photometric redshift. The technique described in Pérez-González et al. (2005) has been improved by significantly increasing the resolution of the templates. This is achieved by fitting the SEDs from the UV to the FIR to stellar population synthesis and dust emission models. With this improvement, by comparing with all the available spectroscopic redshifts, our photometric redshifts have a value of $\Delta(z)/(1+z) < 0.1$ for 88% of the galaxies in the CDF-N and CDF-S, and $\Delta(z)/(1+z) < 0.2$ for 96% of these sources. The average (median) value of $\Delta(z)/(1+z)$ is 0.05 (0.03). Further details about the photometric redshift technique can be found in P. G. Pérez-González et al. (2007, in preparation).

Using this technique, we estimated redshifts for an additional 20 sources, bringing the total number of power-law




FIG. 2.— Relationship between the observed R-band magnitude and hard (2–8 keV) hard X-ray fluxes of power-law galaxies detected in the X-ray (filled circles), weakly-detected in the X-ray (open circles), and non-detected in the X-ray (triangles). For non-detected sources, we plot the 5σ upper limits calculated as described in §5.1. The lines and shading represent the regions populated by AGN ($f_X/f_R < 1$), AGN and starbursts ($-2 < \log f_X/f_R < -1$), and quiescent galaxies, starbursts, and LLAGN ($\log f_X/f_R < -2$) (see Barger et al. 2003; Hornschemeier et al. 2003).

galaxies with redshifts or redshift estimates to 43 (69%). We plot in Figure 3 a comparison between the photometric and spectroscopic redshifts of what we will refer to as the ‘comparison sample’. This sample consists of the 1420 IRAC sources that meet the detection cut of $S/N > 6$ in each of the IRAC bands and that have X-ray exposures ≥ 0.5 Ms; the 11 sources in the original power-law sample removed due to bad or blended photometry were excluded. The mean (median) spectroscopic redshift of the comparison sample is $z = 0.74$ (0.64). Nearly half of the sources in the comparison sample have IRAC SEDs that cannot be fit by a power-law. Of the remaining galaxies, 6% and 40% can be fit by red ($\alpha \leq -0.5$) or blue ($\alpha \geq -0.5$) power-laws, respectively. Only 50% of the X-ray sources in the main CDF-N catalog (Alexander et al. 2003) meet the exposure time and IRAC S/N cuts used to define the comparison sample.

We also plot in Figure 3 the redshift comparison for those galaxies that meet the power-law criterion. While the power-law galaxies tend to be the farthest outliers, the agreement between the spectroscopic and photometric redshifts is reasonable, given the difficulty in assigning redshifts to sources with power-law dominated SEDs. Notably, several power-law galaxies at low spectroscopic redshifts have been assigned high redshifts by our photometric code. We take this into consideration during the analysis, and rely on spectroscopic redshifts alone wherever possible. Nonetheless, the success rate in the photometric redshifts (defined by agreement to 2 standard deviations) implies that our total suite of redshifts is $\geq 80\%$ correct.

As shown in Figure 4, the power-law galaxies tend to lie at significantly higher redshift than both the average source from the comparison sample and X-ray-detected members of the comparison sample, a trend that will be investigated further in §5.1.2. The power-law galaxies are comparable in number to the X-ray-detected AGN in the comparison sample at $z > 2$,

although as will be discussed in §5.1, only half of the power-law galaxies are cataloged X-ray sources. Alonso-Herrero et al. (2006) similarly found that the power-law galaxies have a relatively flat redshift distribution (with the exception of a spike at $z = 1.4$), with most sources lying at $z > 1$.

5. MULTIWAVELENGTH PROPERTIES

In the following sections, we discuss the X-ray, infrared, radio, and optical properties of the power-law galaxies. We show that while only 55% of them have cataloged X-ray counterparts, as many as 85% show evidence for faint X-ray emission. These X-ray-detected power-law galaxies make up a significant fraction of the X-ray luminous AGN in the comparison sample. As such, they are detectable to large distances, explaining the high redshifts of the sample.

The power-law galaxies also have a $24 \mu\text{m}$ detection fraction of nearly unity, which we show to be indicative of an intrinsically luminous X-ray population. While only 29% of the sources have radio counterparts, and all but 2 are radio-quiet, nearly all of the power-law galaxies also have radio luminosities or upper limits consistent with that of AGN-dominated sources. It is therefore likely that the power-law galaxies not detected in the X-ray are intrinsically luminous AGN hidden behind high columns of obscuring gas and dust. The optical–MIR SEDs of the sample, which are flatter than the typical radio-quiet (star-formation subtracted) AGN SED (Elvis et al. 1994), also suggest increasing obscuration in the sources not detected in the X-ray. The optical detection fraction in the GOODS field is high (85%), with approximately 50% of the X-ray sources having compact optical counterparts.

5.1. X-ray Emission

Of the 62 power-law galaxies, only 34 (55%) have cataloged X-ray counterparts in the Alexander et al. (2003) catalog of the 2 Ms CDF-N X-ray field⁴. Sources in the Alexander et al. (2003) catalog were chosen using WAVDETECT (Freeman et al. 2002) with a false-positive probability threshold of 1×10^{-7} . For those power-law galaxies without X-ray counterparts, we searched for faint X-ray emission following the procedure outlined in Donley et al. (2005). We restricted this search to sources with off-axis angles of $\theta < 10'$, as sources that lie outside this radius fall in regions of rapidly changing exposure, and often have sky backgrounds that cannot be well-fit by Poisson distributions; 84% (52) of the power-law galaxies met this criterion. Of these, 58% (30) are detected in the Alexander et al. catalog, 19% (10) have at least a 2.5σ detection in one X-ray band and at one encircled-energy radius (EER)⁵, and 13% (7) remain non-detected. Coadding the 7 non-detected power-law galaxies does not lead to a detection. Four additional sources lie too close to a cataloged X-ray source to search for faint emission, and 1 source (CDFN 8362) has an irregular sky background, preventing us from accurately testing for detection. If we assume that, like those sources for which we could test for emission, 85% of the 5 sources with nearby counterparts or irregular sky backgrounds would be detected (either strongly or weakly), then

⁴ Throughout the paper, we use only the main source catalog from Alexander et al. (2003), and do not include the lower-significance X-ray sources detected in the supplemental optically bright X-ray catalog. No power-law galaxies and only 2 of the color-selected sources discussed in §6.2 are detected in the optically bright catalog.

⁵ The 80% EER refers to the radius within which 80% of the energy is found.




FIG. 3.— Comparison between secure spectroscopic redshifts and photometric redshifts for the comparison sample (left) and the power-law galaxy sample (right). We overplot on each panel a line of slope 1.




FIG. 4.— Top: Redshift distributions of the comparison sample (unshaded) and the power-law galaxy sample (shaded). Bottom: Redshift distributions of the X-ray-detected members of the comparison sample (unshaded) and the X-ray-detected (lightly shaded) and non-detected (heavily shaded) members of the power-law galaxy sample. The distributions based only on spectroscopic redshifts are shown on the left; those based on both photometric and spectroscopic redshifts are shown on the right. The lower panels give the power-law fraction, as a function of redshift.




FIG. 5.— X-ray detection fraction as a function of redshift (top left), X-ray exposure time (top right), power-law spectral index α (bottom left), and power-law fit probability P_x (bottom right). Shaded histograms represent the X-ray-cataloged power-law galaxies; clear histograms represent the full power-law sample.

85% of the power-law galaxies show evidence for X-ray emission, while 15% do not. Changing the detection threshold to 3, 4, and 5 σ changes the detected fraction to 77%, 71%, and 65%, respectively.

We list the X-ray properties of the weakly-detected power-law galaxies in Table 2. If a source was not detected to $> 2.5\sigma$ in one or more of the three bands, we list in Table 2 a conservative 2.5σ upper limit, measured by adding any positive source flux to a 2.5σ limit on the local sky background. The 70% EER was used to calculate both the source counts and sky background; no aperture correction was applied. For those power-law galaxies without a cataloged or weak X-ray counterpart, we list in Table 1 5σ upper limits, calculated as described above.

While Alonso-Herrero et al. (2006) found that their sample of power-law galaxies included about 1/3 of the hard X-ray sources with $24\ \mu\text{m}$ detections, our more conservative selection detects only 14% of the X-ray sources in the Alexander et al. (2003) catalog that meet our exposure time cut of 0.5 Ms and that have $24\ \mu\text{m}$ flux densities of $S_{24} > 80\ \mu\text{Jy}$. (Further restricting the comparison to those X-ray sources that also meet our IRAC significance cut of $S/N > 6$ only increases the selection fraction to 17%). If we consider only those X-ray and MIR sources in the CDF-N catalog with $\log L_x (\text{erg s}^{-1}) > 42$ (to rule out sources dominated by star-formation), the X-ray-detected power-law galaxies comprise $\sim 20\%$ of the X-ray

sources (with exposures of > 0.5 Ms) detected at $24\ \mu\text{m}$ (to $S_{24} > 80\ \mu\text{Jy}$). Because our selection criteria were designed to be quite robust, we exclude a number of IRAC color-selected AGN candidates with power-law like emission (see §6). Determining the intrinsic proportion of “power-law” AGN will therefore require a more careful analysis. As mentioned in §4 and shown in Figure 4, the power-law galaxies comprise a significantly higher fraction of the high-redshift X-ray sources.

5.1.1. Dependence on Sample Properties

We plot in Figure 5 the X-ray detection fraction for the power-law galaxies as a function of redshift, X-ray exposure time, power-law slope, and power-law fit probability, where we consider as ‘detected’ only those sources with cataloged X-ray counterparts. For the sources with redshift estimates, X-ray detection appears to be relatively insensitive to redshift. The X-ray detection fraction, however, is higher for those sources with redshift estimates (70%) than for the full power-law sample (55%). The detection fraction as a function of X-ray exposure time is lower than average at the lowest exposures, as expected, but does not increase substantially at the highest exposures. We note that in our analysis of the X-ray detection fraction in §5.1, we considered only those power-law galaxies with off-axis angles of $\theta < 10'$, and therefore excluded 6 of the 10 sources in the lowest two exposure bins. There is significant variation in the detection fraction as

a function of power-law index, with an increase at the steepest indices and a possible decrease at the flattest indices. The power-law fit probability, however, appears to have a minimal effect on the X-ray detection fraction, with only a slight drop towards the highest values of P_χ (recall that $P_\chi = 0.5$, not $P_\chi = 1$, is ideal). These results suggest that the X-ray detection fraction of the power-law sample is relatively insensitive to variations in redshift, X-ray exposure time, and power-law properties, to the limits imposed by our selection criteria.

5.1.2. X-ray Luminosity

We plot in Figure 6 the observed 0.5–8 keV X-ray luminosities of the power-law galaxies and the members of the comparison sample detected in the X-ray catalog. The luminosities of sources selected via the Lacy et al. selection criteria (see §6.2) are also plotted. As was discussed by Alonso-Herrero et al. (2006), power-law galaxies have X-ray luminosities typical of AGN ($\log L_x(\text{ergs s}^{-1}) \geq 42$), and significantly higher than those of star-forming galaxies ($\log L_x(\text{ergs s}^{-1}) < 41$).

The power-law galaxies comprise a significant fraction of the high luminosity sample. This is not surprising, as the power-law selection requires the AGN to be energetically dominant, and therefore preferentially selects the most luminous AGN. To illustrate the effect of X-ray luminosity on the AGN contribution to the optical through MIR continuum, we plot in Figure 7 the median rest- and observed-frame optical-MIR SEDs of the X-ray-detected members of the comparison sample, as a function of X-ray luminosity. We normalize the SEDs to the $12 \mu\text{m}$ monochromatic power, which, for the rest-frame comparison, is representative of bolometric luminosity and independent of galaxy or AGN type (Spinoglio & Malkan 1989). As can be seen, low-luminosity X-ray sources are dominated by the $1.6 \mu\text{m}$ stellar bump in the optical-NIR bands (e.g. Alonso-Herrero et al. 2004). The relative strength of this feature decreases with increasing X-ray luminosity, and disappears almost entirely at luminosities of $L_x > 44.0$, taking on the characteristic power-law shape required by the selection used here. The fraction of X-ray sources in the comparison sample that can be fit by red, AGN-dominated power-laws ($\alpha \leq -0.5$) is therefore a strong function of X-ray luminosity, increasing from 0% at $\log L_x(\text{ergs s}^{-1}) = 41 - 42$ to 3%, 24%, and 59% at luminosity limits of $\log L_x(\text{ergs s}^{-1}) = 42 - 43$, $43 - 44$, and $44 - 45$. Blue power-laws ($\alpha > -0.5$) in the IRAC bands are indicative of stellar-photosphere-dominated SEDs in the rest-frame near-infrared (for $z \leq 1.5$). The corresponding percentages of X-ray sources fit by blue power-laws are 39%, 46%, 41%, and 14%, showing the reduced proportion of stellar-dominated objects in the highest luminosity bin.

A similar trend is seen as a function of redshift. In a flux limited sample such as this, low-luminosity AGN are detected primarily at low redshift, whereas high-luminosity AGN are detectable over a range of redshifts (see Figure 8). To first order, the high redshifts of the power-law sample can therefore be explained by their preferentially high luminosities. In addition, AGN with high X-ray luminosities have comoving space densities that peak at higher redshifts than low-luminosity AGN, amplifying the selection effect discussed above (e.g. Ueda et al. 2003). The color-redshift relation for AGN may also play a minor role in this trend. Richards et al. (2006) show that the $3.6 \mu\text{m}$ to $5.8 \mu\text{m}$ color ($\log(S_{5.8}/S_{3.6})$) of an AGN with the median radio-quiet SED (Elvis et al. 1994) shifts from 0.2 at $z = 0$ to 0.44 at $z = 1.5$ before falling to 0.08




FIG. 6.— Comparison between the observed 0.5–8 keV X-ray luminosity (L_x) distributions of the cataloged X-ray sources in the comparison sample (unshaded histogram), the power-law sample (filled histogram), and the Lacy et al. sample (lightly shaded histogram; see §6.2). The lower panel gives the fraction of the X-ray sources that meet the power-law or Lacy et al. criteria.

at $z = 4$. As will be shown in §5.4, however, the power-law galaxies tend to have flatter SEDs than the Elvis et al. (1994) AGN template and as such, they should have colors less sensitive to changes in redshift.

5.2. $24 \mu\text{m}$ Emission

Unlike Alonso-Herrero et al. (2006), we do not require a $24 \mu\text{m}$ detection for source selection. However, all but 5 of the power-law galaxies have a cataloged $24 \mu\text{m}$ counterpart within a $2''$ radius. The $24 \mu\text{m}$ flux densities are listed in Table 1. Of the remaining 5 sources, 3 (CDFN 7514, CDFN 14642, and CDFN 25218) have a cataloged counterpart within $2-3''$. For these power-law galaxies, we use the flux density of the nearby source as an upper limit if it exceeds our nominal limit of $80 \mu\text{Jy}$. One additional power-law galaxy, CDFN 45610, lies within the PSF of a bright $24 \mu\text{m}$ source, preventing a detailed analysis of its properties. Only one source, CDFN 38580, an X-ray-detected power-law galaxy at $z = 2.91$ (Chapman et al. 2005), shows no evidence for $24 \mu\text{m}$ emission. As was found for the power-law X-ray sources in the EGS (Barmby et al. 2006), there is only weak agreement between the $24 \mu\text{m}$ flux densities and those predicted by extending the IRAC power-law fit, with the X-ray-detected and X-ray non-detected power-law galaxies having observed $24 \mu\text{m}$ flux densities that tend to lie within only 53% and 60% of the predicted values, respectively.

The fraction of AGN in MIR samples has been shown to increase with increasing $24 \mu\text{m}$ flux density (Treister et al. 2005; Brand et al. 2006) in general agreement with the models of Pearson (2005). We would therefore expect luminous AGN in a flux-limited sample to have a high $24 \mu\text{m}$ detection fraction, such as that seen for the power-law sample. To test this, we plot in Figure 9 the $24 \mu\text{m}$ detection fraction for the members of the comparison sample as a function of X-ray luminosity. The detection fraction is relatively high ($\sim 90\%$) at X-ray luminosities typical of starburst galaxies ($\log L_x(\text{ergs s}^{-1}) < 41$, see Alexander et al. 2002). This is not surprising, as rapidly star-forming galaxies with large numbers of X-ray binaries will also be luminous infrared sources. The detection fraction then drops to $\sim$




FIG. 7.— Median rest-frame (left) and observed-frame (right) SEDs of the X-ray-detected members of the comparison sample, as a function of observed 0.5–8 keV luminosity (in units of ergs s^{-1}). Stars indicate the wavelengths of the four IRAC bands.

65% at luminosities typical of starbursts and low-luminosity AGN ($41 < \log L_x (\text{ergs s}^{-1}) < 42$) before rising to 100% at $\log L_x (\text{ergs s}^{-1}) > 44$ –45. Because the comparison sample is composed of objects with high significance IRAC MIR detections, the $24 \mu\text{m}$ detection fraction is higher than that of an unbiased sample. For comparison, Rigby et al. (2004) find an overall $24 \mu\text{m}$ detection fraction of only 60% for hard X-ray sources in the CDF-S. Nevertheless, for AGN-dominated sources, a high $24 \mu\text{m}$ detection fraction like that of the power-law galaxies accompanies an X-ray-luminous AGN population. We note that of the 18 power-law galaxies not detected in X-rays, all but one are detected at $24 \mu\text{m}$, with the exception being CDFN 45610, the power-law galaxy that lies in the PSF of a bright $24 \mu\text{m}$ source. Therefore, if these sources are AGN-dominated as expected, it is likely that they, along with the rest of the power-law sample (§5.1.2 and Figure 6), are intrinsically X-ray luminous but heavily obscured.

5.3. Radio Loudness

Eighteen of the 62 power-law galaxies (29%) have cataloged radio counterparts from the Richards (2000) 1.4 GHz VLA survey of the CDF-N. This survey has a detection limit of $40 \mu\text{Jy}$ and is 95% complete at $80 \mu\text{Jy}$. In addition, 3 of these power-law galaxies (CDFN 16796, CDFN 27641, and CDFN 37999) are members of the radio-selected submillimeter SCUBA sample of Chapman et al. (2005), as is one radio source that fell below the Richards (2000) detection limit (CDFN 38580). For comparison, Polletta et al. (2006) found that half of their obscured AGN sample were radio-detected, although their radio limits were deeper and their IRAC coverage more shallow.

Since all of the radio-detected power-law galaxies are also detected at $24 \mu\text{m}$, we can test for radio-loudness via the parameter q , where $q = \log (f_{24 \mu\text{m}}/f_{1.4 \text{ GHz}})$ (see Appleton et al. 2004). Sources with $q < 0$ have far more radio emission (with respect to MIR emission) than is expected from the radio/infrared correlation of radio-quiet AGN and star-forming galaxies, and are referred to as radio-excess AGN (e.g. Yun, Reddy, & Condon 2001, Donley et al. 2005). Kuraszekiewicz et al. (2007) show that q is less sensitive to reddening and

host galaxy dilution than measures of radio loudness based on radio and optical flux densities. Only 2 of the 18 radio-detected power-law galaxies are radio-excess AGN; the remaining galaxies are consistent with the radio/infrared correlation⁶. Kuraszekiewicz et al. (2007) find that $\lesssim 15\%$ of AGN are radio-intermediate or radio-loud based on a cut of $q < 0$, consistent with the radio-loud fractions of Kellermann et al. (1989) and Stern et al. (2000). The fraction of radio-excess AGN in the power-law sample ($\sim 3\%$) may therefore be lower than average.

Both sources classified as radio-excess AGN have a cataloged X-ray counterpart; 11 of the remaining 16 radio sources are also bright X-ray sources, while 3 are weakly-detected. Only two radio sources (CDFN 40913 and CDFN 51788) are not detected in the X-ray. As such, selection criteria that include a radio flux density cut may be more likely to select X-ray bright AGN than selection via MIR properties alone. If such a cut were based on the current sensitivity of the CDF-N radio and X-ray surveys, however, only 13 of the 34 power-law galaxies detected in the X-ray catalog (38%) would be selected; the remaining 62% would be missed.

While the majority of the power-law galaxies with radio counterparts appear to be radio-quiet as defined by the radio/infrared correlation, their radio luminosities are consistent with those of AGN-powered sources. In the local universe, galaxies with $\log L_{\text{radio}} (\text{W Hz}^{-1}) > 23$ tend to be AGN-dominated (Condon et al. 2002; Yun et al. 2001). In the GOODS-North (GOODS-N) field, Morrison et al. (2006) show that galaxies with $\log L_{\text{radio}} (\text{W Hz}^{-1}) > 24$ and $z > 1$ have up to an 80% X-ray detection rate. While a source of this luminosity could be a star-formation powered ULIRG (from the radio/infrared correlation, a radio luminosity of $\log L_{\text{radio}} (\text{W Hz}^{-1}) = 24$ corresponds to a total infrared (TIR, $8\text{--}1000 \mu\text{m}$) luminosity of $4 \times 10^{12} L_{\odot}$ (Bell 2003)), the high X-ray detection rate suggests that the population is dominated

⁶ We note that the radio-excess AGN CDFN 12939 was not included in the sample of Donley et al. (2005) because its X-ray exposure time fell slightly below the more stringent cut of 1 Ms. The source CDFN 20246 corresponds to VLA 123709+622258 in Donley et al. (2005).




FIG. 8.— X-ray luminosity vs. redshift for the X-ray sources in the comparison sample (open circles) and the power-law galaxies with available redshift estimates (filled circles). The on-axis X-ray flux limit of $7 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ (Alexander et al. 2003) is plotted for reference.

by AGN.

All 16 of the 18 radio-detected power-law galaxies with redshifts have observed 1.4 GHz luminosities in excess of $\log L_{1.4\text{GHz}} (\text{W Hz}^{-1}) > 23$, and 14 (88%) have $\log L_{1.4\text{GHz}} (\text{W Hz}^{-1}) > 24$. In addition, 26 of the 27 radio non-detected power-law galaxies with redshifts have upper limits greater than $\log L_{1.4\text{GHz}} (\text{W Hz}^{-1}) = 23$ (the exception being CDFN 49937, a low-luminosity X-ray source), and 21 (78%) have limits greater than $\log L_{1.4\text{GHz}} (\text{W Hz}^{-1}) = 24$. Using a cut of $\log L_{1.4\text{GHz}} (\text{W Hz}^{-1}) > 24$, Morrison et al. (2006) found that 60% of luminous radio galaxies in the GOODS-N field have power-law dominant SEDs. Using the same cut, we find that as many as 81% of power-law selected AGN are radio luminous, though not necessarily radio-loud.

5.4. Optical Morphology and Detection Fraction

While it is difficult to determine optical morphologies for faint sources, particularly those with only ground-based optical data, at least 20 of the power-law galaxies have point-like optical/NIR counterparts (8 of which were detected in the GOODS *Hubble Space Telescope* (HST) field, and 12 of which have only ground-based data). Of these, 18 are detected in the X-ray, one is weakly-detected, and one is non-detected. As such, at least 50% of the X-ray-detected sources have optical counterparts dominated by the central engine. Of the 15 sources with BLAGN spectra, 13 are point-like, compared to 2 of the 4 sources with NLAGN spectra. Of the power-law galaxies with available GOODS HST data, $\sim 30\%$ of the sources with optical counterparts are point-like.

Worsley et al. (2006) stacked the X-ray emission from optical sources in the GOODS Fields to test whether these galaxies could account for the remaining unresolved fraction of the CXRB. While the stacked emission of galaxies not individually detected in the X-ray accounts for the unresolved portion of the CXRB at 0.5–6 keV, the reduced sensitivity of *Chandra* at higher energies makes a determination of the resolved fraction more difficult. They estimate that GOODS optical sources account for at most 40% of the unresolved




FIG. 9.— 24 μm detection fraction as a function of X-ray luminosity for the members of the comparison sample detected in the X-ray. The solid and dashed lines give the detection fraction for all sources with redshift estimates and for those with spectroscopic redshifts, respectively. Dotted lines give the upper and lower limit on the 24 μm detection fraction of the power-law galaxies.

emission from 6–8 keV, suggesting that some of the missing CXRB sources may also be missed in the GOODS images. Of the 27 power-law galaxies that lie in the GOODS region, 23 (85%) have cataloged GOODS optical counterparts. Two of the sources missed in the catalog (CDFN 32673 and CDFN 39529) are obscured X-ray sources similar to the extreme X-ray/optical sources (EXOs, e.g. Koekemoer et al. 2004), one (CDFN 33052) is not detected in the X-ray, and one (CDFN 25218) is an obscured, weakly-detected X-ray source that may have a faint B-band counterpart. Therefore, at most 7% (2/27) of the power-law galaxies in the GOODS region are obscured AGN not detected in either the GOODS optical or the X-ray catalogs.

5.5. Optical–MIR SEDs

We plot in Figure 10 the median optical through MIR rest-frame SEDs of the cataloged, weakly-detected, and X-ray non-detected power-law galaxies, normalized by their 1.25 μm monochromatic flux densities. For comparison, we plot the median radio-quiet, starlight-subtracted AGN SED of Elvis et al. (1994). The median SEDs of the power-law galaxies are generally flatter than the Elvis et al. SED, and show no evidence of a UV bump. The median SED of the power-law galaxies with X-ray detections resembles that of the BLAGN SED class of Alonso-Herrero et al. (2006); that of the non-detected power-law galaxies drops much more rapidly in the optical, presumably due to increasing optical obscuration, and resembles that of the the NLAGN SED class of Alonso-Herrero et al. (2006). The weakly-detected sources have optical SEDs that fall between those of the detected and non-detected sources, suggesting an intermediate level of obscuration. The steepest IRAC spectral index found in our sample is $\alpha = -3.15$, only slightly steeper than that found by Alonso-Herrero et al. (2006) ($\alpha = -2.8$) and Rigby et al. (2005) ($\alpha = -2.9$).

Brand et al. (2006) show that X-ray detected AGN in




FIG. 10.— Median optical through MIR rest-frame SEDs of the X-ray-detected (solid line), weakly-detected (dashed line), and non-detected (dotted line) power-law galaxies. The median radio-quiet SED of Elvis et al. (1994) is shown by a solid line that spans the frequency range of the plot. The SEDs have been normalized to their $1.25\ \mu\text{m}$ flux densities.

the 5 ks XBoötes field and MIR-detected sources with clear signs of AGN activity in their optical spectra tend to have flat MIR spectral slopes ($\log \nu f_\nu(24)/\nu f_\nu(8) = 0$, e.g. Elvis et al. 1994), whereas star-forming galaxies typically have higher values of $\log \nu f_\nu(24)/\nu f_\nu(8) \sim 0.5$. Power-law galaxies have $\log \nu f_\nu(24)/\nu f_\nu(8) = -0.24$ to 0.73 , with a mean slope in νf_ν of 0.18 ± 0.22 , suggesting that, when compared to unobscured AGN flat in $\log \nu f_\nu$, either relatively more flux is observed at $24\ \mu\text{m}$ or relatively less flux is observed at $8\ \mu\text{m}$. The power-law galaxies in the X-ray catalog tend to have slightly flatter slopes (0.12 ± 0.21) than those not strongly detected in X-rays (0.26 ± 0.21). This trend can be seen in Figure 10, where the power-law galaxies not detected in the X-ray rise more prominently in $\log \nu f_\nu$ with increasing wavelength than do those galaxies detected in X-rays, suggesting that the X-ray non-detected power-law galaxies are more heavily obscured in the optical.

6. COMPARISON WITH OTHER MIR SELECTION TECHNIQUES

In the following section, we compare the power-law selection technique to a number of other MIR-based AGN selection criteria. While the power-law galaxies display a large range of infrared-to-optical ratios, and cannot be selected on this basis, they comprise a significant fraction of the sources with high infrared-to-optical ratios, suggesting that many heavily optically-reddened sources have power-law continua in the NIR/MIR. The power-law selection more closely matches the *Spitzer* color-color selection criteria of Lacy et al. (2004) and Stern et al. (2005). We compare the X-ray completeness of these techniques, and show that, like the power-law selection, these color criteria tend to select high-redshift X-ray luminous AGN in the deep X-ray fields. We also discuss the X-ray detection fractions of the color-selected samples, which are lower than that of the power-law sample. We argue that this is due, at least in part, to a larger contamination by star-forming galaxies.

6.1. Optical vs $24\ \mu\text{m}$ Emission

The selection of infrared-bright, optically-faint sources has been suggested as a means of identifying obscured AGN (Houck et al. 2005; Weedman et al. 2006; Yan et al. 2004, 2005). Approximately 40 such sources have been observed with the *Spitzer* IRS; the majority (75%) are dominated by

silicate absorption, indicative of heavy obscuration, or are featureless. Although no deep X-ray information is available for those sources observed thus far, their spectral characteristics suggest that many are likely to be obscured AGN.

Only one member of the power-law sample, CDFN 27360 (an X-ray-detected AGN with a column density of $N_H = 2.5 \times 10^{23}\ \text{cm}^{-2}$ (see Table 3)), meets the strict criterion used by Houck et al. (2004) and Weedman et al. (2006) to select infrared-bright, optically-faint sources: $f_{24} > 0.75\ \text{mJy}$ and $R > 24.5$ (Houck et al. 2004), or $f_{24} > 1.0\ \text{mJy}$ and $R > 23.9$ (Weedman et al. 2006). These selection criteria, however, have been set to ensure that sources can be followed-up effectively with IRS. We plot in Figure 11 the positions of the comparison sample (discussed in §4) on the color-color plot of Yan et al. (2004); with the exception of the 7 sources that are X-ray non-detected at the 2.5σ level (see §5.1), 6 of which have no R-band counterpart, only sources detected at $8\ \mu\text{m}$, $24\ \mu\text{m}$, and R-band are shown.

Following Yan et al. (2004, 2005) we define the following quantities:

$$R(24, 0.7) = \log(\nu f_\nu(24\ \mu\text{m})/\nu f_\nu(R)) \quad (1)$$

$$R(24, 8) = \log(\nu f_\nu(24\ \mu\text{m})/\nu f_\nu(8\ \mu\text{m})) \quad (2)$$

According to Yan et al. (2004), sources with $R(24, 0.7) > 1.5$ and $R(24, 8) \sim 0.5$ are likely to be dust-reddened AGN, although 75% of the luminous starburst candidates with $R(24, 0.7) > 1$ and $R(24, 8) > 0.3$ followed-up with IRS by Yan et al. (2005) appear to be either unobscured AGN or galaxies with both a buried AGN and a starburst component. While the power-law galaxies (and X-ray-detected members of the comparison sample) cover a large range of $R(24, 0.7)$ and $R(24, 8)$, as is expected for a sample of AGN with a variety of redshifts and obscurations, the power-law galaxies comprise an increasingly significant fraction of the highly optically-reddened members of the comparison sample with $R(24, 0.7) \geq 1.2$. This suggests that power-law selection is capable of detecting both optically obscured and unobscured AGN, as expected, and that a significant fraction (20-40%) of the infrared-bright/optically-faint sources in the comparison sample have power-law SEDs in the NIR/MIR. In addition, those power-law galaxies shown to be X-ray undetected to the 2.5σ level all have upper limits of $R(24, 0.7) \geq 1$, consistent




FIG. 11.— Location of the comparison sample on the color-color diagram of Yan et al. (2004). All members of the comparison sample are shown (dots), as are the power-law galaxies (diamonds) and the X-ray-detected members of the comparison sample (crosses). The 7 power-law galaxies that are X-ray non-detected to the 2.5σ level are indicated by filled diamonds. Dotted lines show the selection criteria of Yan et al. (2004), as discussed in §6.1. The lower panel shows the power-law fraction of the comparison sample as a function of $R(24,0.7)$.

with the expectations for obscured AGN. In the following section, we compare the power-law selection to selection criteria that more closely match those used here.

6.2. Spitzer Color-Color Selection

MIR AGN color selection criteria have been defined by Ivison et al. (2004), Lacy et al. (2004), Stern et al. (2005), and Hatziminaoglou et al. (2005). The completeness and reliability of the latter three criteria in selecting X-ray-detected AGN in the Extended Groth Strip (EGS) is discussed by Barmby et al. (2005). We plot in Figures 12 and 13 the position of the power-law galaxies with respect to the IRAC color-color cuts of Lacy et al. (2004) and Stern et al. (2005), both of which have been designed for relatively shallow surveys. In addition, we overplot the redshifted IRAC colors of the Dale & Helou (2002) star-forming template (which is degenerate with their parameter α for the wavelengths of interest), the median radio-quiet AGN SED of Elvis et al. (1994), and the SEDs of the ULIRGs Mrk 273, IRAS 17208-0014, and Arp 220, described in Appendix A.

The selection criteria of Lacy et al. are based on SDSS quasars, and therefore are not designed to select AGN in which the host galaxy dominates the MIR energy, as well as AGN that are obscured in the MIR. As discussed in Alonso-Herrero et al. (2006) and as shown in Figure 12, IRAC power-law galaxies fall along a line well within the Lacy et al. selection region, although these galaxies fill only a small fraction of the available color space within the defined cut. Only 16% of the objects from the comparison sample that satisfy the Lacy et al. color cuts also meet the power-law criteria.

The Stern et al. (2005) selection criteria provide a slightly closer match to the power-law selection technique. Stern et al. define their color cut using ~ 9400 sources from AGES⁷,

800 of which are spectroscopically confirmed AGN. Their color criterion identifies 91% of the BLAGN and 40% of the NLAGN in the AGES sample. 17% of the sources that met their AGN criteria were not classified as AGN based on their optical spectra, but may be optically normal (optically dull) AGN. All of the power-law galaxies meet the Stern et al. criteria, and 29% of the sources from the sample that meet the Stern et al. criteria also meet the power-law criteria. The power-law galaxies show a larger scatter about the power-law locus in the Stern et al. diagram (Figure 13) than they do in the Lacy et al. diagram (Figure 12); this is likely to be due, at least in part, to the smaller wavelength baseline probed by the Stern et al. colors.

6.2.1. X-ray completeness of color-selected samples

In the EGS, with an X-ray exposure time of ≤ 200 ks, the Lacy et al. and Stern et al. color criteria selected 73% and 51% of the X-ray AGN, respectively (Barmby et al. 2005). In the deeper CDF-N field, these criteria select only 21% and 17% of the Alexander et al. (2003) X-ray sample. However, only 50% of the X-ray sources in the CDF-N meet the exposure time and IRAC S/N cuts used to define the comparison and color-selected samples. When we consider only those X-ray sources in the comparison sample, the Lacy et al. and Stern et al. criteria select 39% and 33% of the X-ray sources, respectively. Further restricting the comparison to sources in the comparison sample with X-ray luminosities indicative of AGN activity, $\log L_x \text{ (erg s}^{-1}\text{)} \geq 42$, increases the selection fractions to 52% and 47%, respectively. For comparison, our strict power-law criteria recover $\sim 20\%$ of the X-ray luminous AGN in the comparison sample.

Power-law galaxies, a subsample of both the Lacy et al. and Stern et al. criteria, tend to preferentially lie at both high luminosity and high redshift. The same is true for the full color-selected samples, as shown in Figures 14 and 15, in which we plot only sources with spectroscopic redshifts and only the Lacy et al. color criteria. Including photometric redshifts does not change the results, and the same trends are seen for both the Lacy et al. and Stern et al. selection techniques.

At low redshift, only a small fraction of the comparison sample lies inside the Lacy et al. selection region (see Figure 14). At redshifts of $z > 1$, however, the vertical branch in color-space populated by low-redshift ($z < 0.3$) aromatic-dominated sources (Sajina, Lacy, & Scott 2005) disappears, and at $z > 2$, all of the remaining IRAC sources meet the Lacy et al. criteria. Similarly, at $\log L_x \text{ (erg s}^{-1}\text{)} < 42$, only a small fraction of the sources in the comparison sample fall in the selection region (see Figure 15). In contrast, 72% of the X-ray sources in the comparison sample with $\log L_x \text{ (erg s}^{-1}\text{)} > 43$ and 100% of those sources with $\log L_x \text{ (erg s}^{-1}\text{)} > 44$ meet the Lacy et al. selection criteria. These high luminosity sources lie in the same region of color space as the power-law selected AGN (see Figure 12), although as is apparent from Figure 6, the Lacy et al. criteria (and the Stern et al. criteria) select a more complete sample of AGN with high X-ray luminosities than does the power-law selection. As is also shown in Figure 6, however, the X-ray luminosities of the color-selected sources extend below the traditional AGN limit of $\log L_x \text{ (erg s}^{-1}\text{)} > 42$, suggesting either heavy obscuration of highly-luminous AGN, or possible contamination by star-forming galaxies.

⁷ the AGN and Galaxy Evolution Survey, C. S. Kochanek et al., in prep




FIG. 12.— Location of the comparison sample on the color-color diagram of Lacy et al. (2004). Symbols are as described in Figure 11. Overplotted are the power-law locus from $\alpha = -0.5$ to $\alpha = -3$ (thin black line) and the redshifted IRAC colors of a typical star-forming galaxy (Dale & Helou 2002, cyan, star), the cold (starburst dominated) ULIRGs Arp 220 (blue, triangle) and IRAS 17208-0014 (green, upside down triangle), the ULIRG/Se2 Mrk 273 (red, square) and the radio-quiet AGN SED from Elvis et al. (1994, magenta, circle), where the indicated point represents the colors at $z = 0$, and small circles mark the colors at $z = 1, 2$, and 3 .

6.2.2. AGN Reliability

Of the sources selected via the power-law criteria, 55% were detected in the X-ray catalog of Alexander et al. (2003). The X-ray detection fractions of the sources selected via the Lacy et al. and Stern et al. criteria are lower: 25% and 38%, respectively. We searched for faint X-ray emission from the color-selected sources using the method described in §5.1; 65% of the Lacy et al. sources and 70% of the Stern et al. sources show evidence for X-ray emission (at the 2.5σ level or higher), compared to 85% for the power-law galaxies. Do these techniques select a larger fraction of AGN not detected in X-rays, or do they suffer from greater contamination from star-forming galaxies?

A spectroscopic follow-up of candidate obscured AGN selected via the Lacy et al. criteria reveals that most ($\sim 80\%$) have spectral signatures typical of type-2 AGN (Lacy et al. 2005). However, the 12 candidate sources for which spectra were taken were chosen to have $8\ \mu\text{m}$ flux densities $> 1\ \text{mJy}$, and all lie at $z < 1.34$. As shown in Figure 12, the SEDs of the star-formation dominated ULIRG Arp 220 enters the Lacy et al. selection region at low ($z = 0.23$ to 0.67) and moderately high ($z > 1.43$) redshift, as does that of IRAS 17208-0014 ($z = 0.36$ to 0.89 and $z > 1.35$). The Dale & Helou (2002) star-forming template shows similar behavior, falling just inside the AGN selection region at low redshift ($z = 0.48$ to 0.58) and re-entering the region at high redshift ($z > 1.66$), suggesting that pure color-selection is capable of selecting both normal and highly-luminous star-forming galaxies at low and moderately high redshift. The simulations of Sajina et al. (2005) also show a significant number of aromatic feature-dominated sources near the outskirts of the Lacy et al. AGN selection region, and suggest that while the IRAC colors provide an ef-




FIG. 13.— Location of the comparison sample on the color-color diagram of Stern et al. (2005). Symbols are as described in Figure 11 and templates are as described in Figure 12.

fective means of identifying obscured AGN up to $z = 2$, such criteria are less effective at higher redshift, where the IRAC bands sample the NIR light from the galaxy (see also Alonso-Herrero et al. 2006). As expected, the AGN template (Elvis et al. 1994) lies in the same region of color space as the power-law galaxies, as does the low-redshift template of Mrk 273, a ULIRG with AGN signatures in the optical and NIR (Veilleux et al. 1999; Risaliti et al. 2006). Mrk 273 meets the Lacy et al. criteria at all redshifts except $z = 0.83$ to 1.40 .

The Stern et al. criteria also suffer from potential contamination from cold ULIRGs and star-forming galaxies, although the star-forming templates tend to fall in the selection region for smaller redshift intervals than in the Lacy et al. color-space. The star-forming template of Dale & Helou (2002) lies along the edge of the selection region at low redshift ($z = 0.51$ to 0.56) but it does not re-enter the selection region at the redshifts plotted here ($z < 3$). Arp 220 falls in the selection region twice (from redshifts of $z = 0.28$ to 0.55 and $z = 1.17$ to 1.44), as does IRAS 17208-0014 ($z = 0.49$ to 0.58 and $z = 1.15$ to 1.60). Once again, both the AGN template (Elvis et al. 1994) and the low-redshift Mrk 273 template have colors similar to those of the power-law galaxies. The Mrk 273 template meets the Stern et al. criteria in three redshift intervals ($z = 0$ to 0.06 , $z = 0.19$ to 0.39 , and $z = 1.22$ to 1.59), whereas the AGN template falls in the selection region at all redshifts plotted here.

The Mrk 273, Arp 220, and IRAS 17208-0014 templates all lie near regions of color-space populated by power-law galaxies. The template of Mrk 273, a ULIRG with AGN signatures, meets our power-law criteria ($\alpha \leq -0.5$, $P > 0.1$, monotonically rising) at redshifts of $z = 0$ to 0.08 , $z = 0.18$ to 0.39 and at $z > 2.88$ (assuming 10% flux errors in the 3.6 and $4.5\ \mu\text{m}$ bands, and 15% flux errors in the 5.8 and $8.0\ \mu\text{m}$ bands). The Arp 220 template, however, does not meet these criteria at redshifts less than $z \sim 2.9$, as expected. With the exception of a narrow redshift window of $z = 0.51$ to 0.53 in which IRAS 17208-0014 appears as a power-law galaxy, the




FIG. 14.— Location in the Lacy et al. (2004) color space of the comparison sample as a function of redshift. Only sources with spectroscopic redshifts are shown. X-ray sources in the comparison sample are indicated by crosses.




FIG. 15.— Location in the Lacy et al. (2004) color space of the comparison sample as a function of X-ray luminosity (in units of ergs s^{-1}). Only sources with spectroscopic redshifts are shown.




FIG. 16.— Total infrared luminosity (8-1000 μm) required for Arp 220 to meet the detection limits of our survey (see §3). The lines represent the luminosity required for detection in the MIPS 24 μm (solid) and IRAC 3.6 μm (long-dashed), 4.5 μm (short-dashed), 5.8 μm (dotted), and 8.0 μm (dot-dashed) bands.

same is true for this star-forming template, which meets our criteria at $z \gtrsim 2.8$. Only five of the sources in our sample with redshift estimates lie at $z < 1$; none of these lie between $z = 0.51$ and 0.53 , and all are detected in the X-ray. We therefore expect little contamination from low-redshift ULIRGs, although the presence of such sources cannot be ruled out. As for contamination from high redshift star-forming galaxies, a ULIRG with an SED like that of Arp 220 that lies at $z = 2.8$ must have a total infrared luminosity (TIR, 8-1000 μm) of $\log L_{\text{TIR}}(L_{\odot}) > 13.3$ to have a 24 μm flux density $> 80 \mu\text{Jy}$ (see Figure 16). The fraction of ULIRGs dominated by AGN increases with infrared luminosity, with 50% of ULIRGs showing Seyfert-like properties at $\log L_{\text{TIR}}(L_{\odot}) > 12.3$ (Veilleux et al. 1999). It is therefore unlikely that HyperLIRGs with $\log L_{\text{TIR}}(L_{\odot}) > 13$ have NIR luminosities dominated by star-formation. As such, we also expect little contamination in the power-law sample from high-redshift star-forming galaxies, so long as the power-law objects are also detected at 24 μm (as is the case both for the sample of Alonso-Herrero et al. (2006) and for the majority of our sample).

As a final check, we plot in Figures 12 and 13 the seven power-law galaxies in our sample that are not detected in the X-ray at the 2.5σ level. With one or two exceptions, they are well removed from the colors of Arp 220 and IRAS 17208-0014 for any value of $z < 2.8$ (particularly in Figure 12). We therefore believe they contain AGN that are heavily obscured in the X-ray. This test illustrates the utility of the color-color plots in identifying regions of color space within the color selection regions and near the power-law locus in which contamination by star-forming galaxies is possible. The current templates suggest that pure color selection is likely to select star-forming galaxies and ULIRGs at both low and high redshifts. Several other lines of evidence also point towards a higher fraction of star-forming galaxies in the color-selected samples. While $\geq 92\%$ of the power-law galaxies are detected at 24 μm , the 24 μm detection fractions of the Lacy et al. and Stern et al. samples are lower: 72% and 83%, respec-

tively. In addition, while 88% of the radio-detected power-law galaxies with redshift estimates have radio luminosities of $L(1.4\text{GHz}) > 24 \text{ W Hz}^{-1}$, the value typical of AGN-powered radio galaxies in GOODS-N (Morrison et al. 2006), only 49% and 54% of the Lacy et al. and Stern et al. radio sources (with redshift estimates) have luminosities that exceed this value. Finally, we plot in Figure 17 the position of the color-selected sources on the X-ray to optical diagnostic diagram. While the current detections and upper limits place only two power-law galaxies within the transition region and none near the region populated by quiescent galaxies (see Figure 2), 15% and 10% of the Lacy et al. and Stern et al. sources fall in these regions, and more have upper limits that place them just outside.

6.2.3. SEDs of color-selected sources

How do the MIR SEDs of the Lacy et al. and Stern et al. selected sources compare to those of the power-law galaxies? Of the Lacy et al. sources, 20% can be fit by red, AGN-dominated power-laws ($\alpha < -0.5$) and 30% can be fit by blue, stellar-dominated power-laws ($\alpha > -0.5$). A higher fraction of the Stern et al. sources have good power-law fits: 37% are red power-laws and 34% are blue power-laws⁸. The comparison sample (see Section 4) shows the behavior of objects selected without regard to the presence of an AGN; 6% and 39% of the full comparison sample can be fit by red and blue power-laws, as can 16% and 35% of the cataloged X-ray sources in the comparison sample.

7. OBSCURATION

In the local universe, Seyfert 2's are four times more numerous than Seyfert 1's (Maiolino & Rieke 1995) and at least half of all Seyfert 2's are Compton-thick (Maiolino et al. 1998; Risaliti et al. 1999). In the more distant universe, X-ray background and luminosity function synthesis models predict global obscured ($\log N_{\text{H}}(\text{cm}^{-2}) \geq 22$) ratios of 3:1 to 4:1 (Comastri et al. 2001; Ueda et al. 2003; Gilli 2004; Treister et al. 2005; Tozzi et al. 2006), significantly higher than the observed type 2 to type 1 ratio of spectroscopically identified X-ray sources in the deep fields, 2:1 (e.g., Barger et al. 2003; Szokoly et al. 2004; Treister et al. 2005), but slightly lower than the 6:1 ratio observed for high-redshift X-ray-detected SCUBA galaxies (Alexander et al. 2005).

In the following discussion, we estimate the X-ray column densities of the power-law sample and compare the resulting obscured fraction to predictions from the XRB. If we consider only those power-law galaxies detected in the X-ray, we find an obscured ratio of 2:1. Including the power-law galaxies both weakly- and non-detected in the X-ray results in an obscured ratio of $\lesssim 4:1$. We use a Monte Carlo code to measure the dispersion in this ratio, and investigate the change in the obscured fraction with X-ray luminosity and redshift. By comparing the space density of obscured AGN in the power-law sample to the predictions of Treister et al. (2006), we estimate that at most 20-30% of obscured, MIR-detected AGN have SEDs that meet our robust power-law criteria. We also discuss the effect of large obscuring columns on the NIR/MIR continuum, and show that while heavily obscured AGN can have NIR emission dominated by the AGN,

⁸ We note we have applied only a power-law fit probability to derive the above numbers, and have excluded the other criteria discussed in §3 (i.e. no turnovers in flux density, no apparent stellar bump). The number of galaxies identified with red power-laws using just these two criteria is therefore 79, not 62.




FIG. 17.— Relationship between the observed R-band magnitude and hard (2–8 keV) X-ray fluxes for the Lacy et al. (2004) sources (left) and the Stern et al. (2005) sources (right). Symbols, lines, and shading are as described in Figure 2.

the power-law criteria may be biased against the most heavily obscured (Compton-thick) AGN.

7.1. Column Densities

We estimated the intrinsic column density, N_H , of each X-ray-detected power-law and color-selected galaxy by comparing the observed hard to soft-band X-ray flux ratio to that expected for a typical AGN at a given redshift, after correcting the flux ratios for a Galactic column density of $N_H = 1.6 \times 10^{20} \text{ cm}^{-2}$. When no redshift was available, we estimated the column densities at redshifts of $z = 0.5$ and $z = 3.0$, the approximate limits of the power-law sample. We assumed an intrinsic photon index of $\Gamma = 1.81$ (Tozzi et al. 2006), and did not include a Compton reflection component, which tends to decrease the necessary column by $\sim 20\%$ (see Donley et al. 2005). The estimated column densities of the power-law galaxies are given in Tables 2 and 3 and plotted in Figure 18.

7.1.1. Obscured fraction

A comparison between the X-ray-cataloged and weakly-detected power-law galaxies is complicated, because only one of the latter has both a hard and soft-band detection. We can therefore place only upper limits on the columns of 6 of the weakly-detected AGN, lower limits on two, and no limit on the remaining one. However, all weakly-detected power-law galaxies for which we can estimate N_H are consistent with being obscured ($N_H > 10^{22} \text{ cm}^{-2}$), but not Compton-thick ($N_H > 10^{24} \text{ cm}^{-2}$). Of the X-ray-detected power-law galaxies, $\sim 68\%$ are obscured, in agreement with obscured fractions found by Ueda et al. (2003) for AGN at similar redshifts. If we consider only those power-law galaxies in the X-ray catalog, we therefore derive an obscured to unobscured ratio of $\sim 2:1$.

Several lines of evidence point towards high obscuration in the sources not detected in X-rays. First, the power-law selection criteria (§5.1.2) and high $24 \mu\text{m}$ detection fraction (§5.2) appear to be indicative of a population of intrinsically X-ray luminous sources (see Fig. 10), sources that, if un-

obscured, should be detectable out to high redshift (see Fig. 9). Secondly, the median optical–MIR SED of the sources not detected in X-rays falls below that of the weakly-detected sources at optical wavelengths, which in turn falls below that of the strongly-detected sources, suggesting a continuously increasing optical obscuration for a given $1.25 \mu\text{m}$ flux density (see Fig. 11). Adding the 10 weakly-detected power-law galaxies, all of which are likely to be obscured (see Table 2), we calculate an obscured fraction of $\sim 75\%$. If we further include the power-law galaxies not detected in X-rays, assuming that all are obscured, the maximum obscured fraction of power-law galaxies rises to 82% (4:1-5:1). (Considering only those power-law galaxies with $\theta < 10'$ gives an obscured fraction of 81%). This upper limit on the obscured fraction is consistent with the fraction of NLAGN SEDs found by Alonso-Herrero et al. (2006) for power-law sources in the CDF-S ($\sim 75\%$).

To place a lower limit on the obscured fraction, we consider two additional scenarios. If we assume that all of the X-ray non-detected power-law galaxies are obscured, as before, but assume instead that the 6 weakly-detected power-law galaxies with upper limits on their column densities are *unobscured*, the obscured fraction drops to $\sim 71\%$. If we further consider only those power-law galaxies with column density estimates (i.e. those sources strongly or weakly detected in the X-ray) and assume again that all of the weakly-detected power-law galaxies with upper limits on their column densities are unobscured, the minimum obscured fraction drops to $\sim 60\%$. This lower limit, however, falls below the obscured fraction of the X-ray-detected power-law galaxies (68%). As it is unlikely that the power-law galaxies not detected in the X-ray are significantly less obscured than those detected in the X-ray, the obscured fraction of the X-ray-detected power-law galaxies can be taken as the lower limit on the obscured fraction.

To investigate the dispersion in the observed ratio of obscured to unobscured AGN, we ran a Monte Carlo simulation in which we varied both the assumed intrinsic photon index, Γ , and the photometric redshifts, and then recalculated the column densities of the power-law galaxies. The intrinsic photon index was drawn from the distribution of Tozzi et al (2006), as measured for the brightest 30 sources in their




FIG. 18.— Distribution of X-ray column densities $N_H(\text{cm}^{-2})$ for the power-law galaxies in the X-ray catalog (unshaded) and those only weakly-detected (shaded). We place only upper or lower limits on the column densities of all but one of the weakly-detected sources; the limits of sources with redshift estimates are shown by small arrows. Sources with undetectably low column densities were assigned columns of $\log N_H(\text{cm}^{-2}) = 19$ for plotting.

sample: $\Gamma = 1.81 \pm 0.20$. We allowed the photometric redshifts of the power-law galaxies to vary about their mean by $\Delta(z) = 0.1(1+z)$ (see §4). For the simulation, we assigned the 6 cataloged and weakly-detected sources with hard-to-soft flux ratios but no redshift estimate the median redshift of our sample, $z = 1.9$. The results are shown in Figure 19. The maximum obscured fraction varies from $\sim 75\%$ to 85% (3:1-6:1), with a mean of 81% (4:1).

7.1.2. Redshift and luminosity dependence of obscured fraction

We plot in the right panel of Figure 19 the obscured fraction as a function of redshift and observed X-ray luminosity. Solid and dotted lines represent power-law galaxies with low ($\log L_x(\text{erg s}^{-1}) < 44$) and high ($\log L_x(\text{erg s}^{-1}) > 44$) absorption-corrected X-ray luminosities, and thin and thick lines represent power-law galaxies with low ($z < 2$) and high ($z > 2$) redshifts, respectively. We include all X-ray sources with redshifts for which column densities are available, and exclude those sources with only upper limits on the X-ray luminosity. In addition to having a relatively high obscured to unobscured ratio, power-law galaxies appear to be more heavily obscured at low X-ray luminosities and at high redshifts. The upper limit on the obscured fraction varies from 0.5, for low-redshift, high-luminosity power-law galaxies, to 1, for high-redshift, low-luminosity sources. As we can plot only those power-law galaxies detected in the X-ray for which redshift estimates are available (37/63), systematic effects are likely to be present (e.g. Treister et al. 2005).

The trends we find are consistent with previous results. A number of authors have predicted an increase in the obscured fraction of AGN with increasing redshift (e.g., Gilli et al. 2001, 2003; La Franca et al. 2005; Ballantyne et al. 2006; Tozzi et al. 2006), although a constant obscured fraction also appears to be consistent with the current data (Ueda et al. 2003; Gilli 2004; Treister et al. 2004; Treister & Urry 2005). Akylas et al. (2006) suggest that the ob-

served increase in the obscured fraction with redshift may be due to statistical fluctuations in the small number of counts in the low energy bands, which can cause the column densities of high redshift sources to be overestimated. Stronger evidence exists for a decrease in the obscured fraction with increasing X-ray luminosity (Ueda et al. 2003; Steffen et al. 2003; Hasinger 2004; Szokoly et al. 2004; Barger et al. 2005; Treister et al. 2005; Ballantyne et al. 2006), as expected from the receding torus model (Lawrence 1991). For instance, Szokoly et al. (2004) find X-ray Type 2 fractions of 75%, 44%, and 33% for AGN with X-ray luminosities of $\log L_x(\text{erg s}^{-1}) = 42-43$, $43-44$, and $44-45$, respectively, and Ballantyne et al. (2006) find that the model that simultaneously best describes local AGN data, the CXRB spectrum, and X-ray number counts requires an obscured ratio of 4:1 for low luminosity AGN and 1:1 to 2:1 for AGN with $\log L_x(\text{erg s}^{-1}) > 44$.

7.1.3. Obscured fraction of color-selected samples

If we assume that all of the color-selected sources are AGN, and that all X-ray-non-detected sources are obscured, the Lacy et al. and Stern et al. selection criteria give upper limits on the obscured fraction of 92% (12:1) and 88% (7:1). As discussed in §6.2.2, however, it is likely that these samples contain star-forming galaxies as well as AGN; these estimates of the obscured fractions are therefore almost certainly high. By comparing the numbers of IR color-selected and optically-selected AGN, Stern et al. (2004) estimated an obscured AGN fraction of 76%, similar to that found for the power-law galaxies, although Richards et al. (2006) use a different infrared-to-optical flux ratio to revise this estimate to $\geq 41\%$. Lacy et al. (2004) similarly found an obscured ratio of 46% for the brightest members of their sample. It is important to note, however, that both the color- and power-law criteria select luminous AGN whose NIR/MIR emission is dominated by the central engine. Low-luminosity and heavily-obscured AGN whose NIR/MIR emission is dominated by the host galaxy will tend to fall outside of the color-selection and power-law regions, increasing the total obscured fraction of AGN.

7.2. Space Densities of Obscured AGN

If we assume that all of the power-law galaxies not detected in X-rays are obscured, the power-law sample contains at most ~ 51 obscured AGN and at least ~ 11 unobscured AGN (assuming that CDFN 50281, with a column density of $N_H \leq 10^{22} \text{ cm}^{-2}$ is unobscured). How do these numbers compare to the predicted space densities of Type 1 and Type 2 AGN?

Treister et al. (2006, private comm.) predict 199 AGN in a 350 arcmin^2 area (the area of our survey) down to a $24 \mu\text{m}$ limiting flux density of $80 \mu\text{Jy}$, $\sim 105-110$ of which should be detected in the hard X-ray band and ~ 174 of which are obscured (Type II). For comparison, the X-ray catalog of Alexander et al. (2003) contains 147-149 hard X-ray sources that meet our exposure cut (and therefore lie in the survey area) and which have $S_{24} > 80 \mu\text{Jy}$ and AGN-like X-ray luminosities of $\log L_x(\text{ergs s}^{-1}) > 42$. At a flux density of $80 \mu\text{Jy}$, the predicted number counts (for X-ray detected AGN) therefore lie within a factor of ~ 1.4 of the observed values.

As discussed in §5.2, all but 5 of the power-law galaxies have $24 \mu\text{m}$ counterparts and all but one of the MIPS-detected sources have a $24 \mu\text{m}$ flux density $> 80 \mu\text{Jy}$. Of the 5 sources without $24 \mu\text{m}$ counterparts, however, only one




FIG. 19.— Left: Upper limit on the fraction of obscured ($\log N_H(\text{cm}^{-2}) \geq 22$) power-law galaxies. All power-law galaxies not strongly or weakly detected in X-rays were assumed to be obscured. Right: Upper limit on the obscured fraction as a function of redshift and X-ray luminosity. Sources with absorption-corrected $\log L_X (\text{erg s}^{-1}) < 44$ are shown by solid lines; dotted lines represent sources with $\log L_X (\text{erg s}^{-1}) > 44$. Thin and thick lines represent sources with redshifts of $z < 2$ and $z > 2$, respectively. For viewing clarity, we do not plot the histograms, but instead connect their points.

source is a clear non-detection. The power-law sample therefore contains 56-60 galaxies with $24 \mu\text{m}$ fluxes $> 80 \mu\text{Jy}$. Based on the Treister et al. (2006) predictions, these sources should account for $\sim 28-30\%$ of the MIR-detected AGN (or $\sim 20-22\%$ if we boost the Treister et al. counts by a factor of 1.4). Likewise, the 45-49 obscured power-law galaxies with $24 \mu\text{m}$ fluxes $> 80 \mu\text{Jy}$ should account for at most $\sim 26-28\%$ ($\sim 18-20\%$) of the obscured, Type II, MIR-detected AGN.

Our results therefore suggest that at most $\sim 20-30\%$ of the MIPS-detected AGN and MIPS-detected obscured AGN have high S/N power-law IRAC continua; the remainder could have SEDs dominated by or strongly affected by the host galaxy, red power-law SEDs that fall below our IRAC detection limit, or have been rejected due to noise that caused them to fail our relatively stringent signal-to-noise criteria. It is not surprising that we find a fairly small portion of power-law AGN, as our selection criteria require the AGN to be both of high luminosity and energetically dominant over the other source components in the near- to mid-IR. Franceschini et al. (2005) find that of a sample of *Chandra*-detected AGN in the SWIRE survey, only 62% have optical-IR SEDs typical of Type 1 or Type 2 AGN, and 40-60% of X-ray-selected AGN are known to be optically-dull (Hornschemeier et al. 2001; Barger et al. 2001; Giacconi et al. 2001). The fraction of AGN with optical-IR SEDs dominated by star-formation is likely to be even higher for heavily obscured AGN samples, such as that identified here. Polletta et al. (2006) select obscured AGN candidates on the basis of red non-stellar SEDs, similar to the power-law criterion used here; only 40% of their obscured AGN candidates show AGN signatures in the optical/NIR SEDs.

7.3. Effect of Reddening on Obscured AGN

Is it reasonable to assume that heavily obscured AGN at high redshift could have NIR/MIR SEDs still dominated by the central engine? Observations of QSOs indicate that AGN can be up to 2-3 magnitudes brighter than their host galaxies in the NIR (e.g. McLeod & Rieke 1994; Percival et al. 2001; Marble et al. 2003), although the contribution of

the non-stellar continuum varies from source to source and as a function of wavelength. The total H-band contribution of luminous AGN, for example, ranges from ~ 35 to 90% (McLeod & Rieke 1994), and while the non-stellar continuum of NGC 1068 accounts for $> 80\%$ of the flux at $2.3 \mu\text{m}$ in the inner $4''4$, the contribution drops to 30% at $1.6 \mu\text{m}$ (Origlia et al. 1993).

In the Milky Way, a column density of $\log N_H(\text{cm}^{-2}) = 23$ corresponds to an A_V of ~ 50 mag (Bohlin, Savage, & Drake 1978). The A_V to N_H ratio of AGN, however, appears to be an order of magnitude lower than that of the Milky Way (e.g. Maccacaro, Perola, & Elvis 1982; Reichert et al. 1985; Granato et al. 1997; Maiolino et al. 2001a), due either to a higher gas-to-dust ratio, or to the formation of large grains in the dense AGN environment (Maiolino et al. 2001a, 2001b). Following Martínez-Sansigre (2006), we therefore adopt an AGN N_H to A_V conversion of 2.0×10^{-23} , for which column densities of $\log N_H(\text{cm}^{-2}) = 22, 23$, and 24 correspond to $A_V = 0.2, 2$, and 20 mag, respectively.

Assuming the IR extinction law of Rieke & Lebofsky (1985), we estimate the observed IRAC extinctions for an obscured AGN at $z = 2$, where IRAC samples the $1-3 \mu\text{m}$ rest-frame emission. At a column density of $\log N_H(\text{cm}^{-2}) = 22$, the extinctions at the center of the IRAC channels are negligible ($0.02-0.06$ mag), but they rise with column density to $0.2-0.6$ mag for $\log N_H(\text{cm}^{-2}) = 23$ and $1.8-6.2$ mag for $\log N_H(\text{cm}^{-2}) = 24$. It is therefore reasonable to assume that luminous obscured AGN could dominate the emission of their host galaxies at all but the highest column densities.

If the interstellar extinction law is applicable to AGN, differential extinction across the IRAC bands will redden the observed spectral slope, potentially increasing the number of heavily obscured luminous AGN in the power-law sample. At $z = 2$, a column density of $\log N_H(\text{cm}^{-2}) = 22$ has only a minor effect on the measured slope: an observed spectral index of $\alpha = -0.5$ corresponds to an unreddened intrinsic slope of $\alpha = -0.45$. At higher column densities of $\log N_H(\text{cm}^{-2}) = 23$ and $\log N_H(\text{cm}^{-2}) = 24$, however, the best-fit intrinsic slopes at $z = 2$ increase to $\alpha = 0.0$ and $\alpha = 4.5$. The unrealistic predicted

intrinsic slope of the high-redshift Compton-thick model suggests that highly obscured AGN at high redshift are likely to have steep observed spectral slopes, if their IR and X-ray emission are subject to the same obscuring material. For example, the spectral slope of CDFN 44836, $\alpha = -3.2$, could be caused by the extinction of an $\alpha = 0$ source by a column density of $\log N_{\text{H}}(\text{cm}^{-2}) = 23.8$ at $z = 2$. At such high column densities, however, the de-reddened source begins to steepen at the bluest bands, no longer resembling a power-law source.

While the above results therefore suggest that power-law selection may be biased against Compton-thick AGN, there exist a population of AGN whose X-rays are more heavily obscured than their optical/NIR emission (e.g. Akiyama et al. 2003; Brusa et al. 2003; Page et al. 2003; Wilkes et al. 2005). For instance, 10% of the BLAGN sample of Perola et al. (2004) have X-ray columns of $\log N_{\text{H}}(\text{cm}^{-2}) \geq 22$. Shi et al. (2006) also find several Compton-thick AGN that do not follow an observed correlation between X-ray column density and the strength of the silicate emission or absorption feature. Instead, their silicate absorption is weaker than expected, suggesting a Compton-thick absorber that obscures the X-ray emission, but not the IR emission. These outliers could also be explained, however, by a Compton-thick absorber that obscures the mid-IR so strongly that the output in this range is dominated by star formation in the host galaxies.

8. CONCLUSIONS

We define a sample of 62 power-law galaxies in the CDF-N. Sources were required to be detected to $S/N > 6$ in each of the IRAC bands, have IRAC slopes of $\alpha < -0.5$ (where $f_{\nu} \propto \nu^{\alpha}$), and to lie in regions with both deep X-ray (> 0.5 Ms) and radio coverage. We studied the multiwavelength properties of the power-law galaxies and compared the power-law selection technique to other MIR-based AGN selection criteria. We then measured the intrinsic obscuring column densities of the power-law galaxies to estimate the obscured fraction of the sample, which should be less affected by obscuration than optical and/or X-ray selected samples. The main results of this paper are as follows:

- Power-law selection requires the AGN to be energetically-dominant in the near/mid infrared. On average, AGN with X-ray luminosities of $\log L_{\text{x}} (\text{erg s}^{-1}) < 44$ have SEDs dominated by or strongly affected by the stellar continuum. Therefore, power-law galaxies tend to have high X-ray luminosities, and make up a significant fraction of the X-ray luminous AGN population.
- Power-law galaxies lie at significantly higher redshifts than the typical IRAC- or X-ray-detected source, primarily due to their high luminosities. While the power-law sample defined here accounts for only $\sim 20\%$ of the X-ray and MIR-detected AGN in the comparison sample, approximately 50% of the high-redshift X-ray sample are power-law galaxies.
- 45% of the power-law galaxies are not detected in the X-ray catalog of Alexander et al. (2003) at exposures of > 0.5 Ms. A search for faint emission reveals that 15% remain undetected at the 2.5σ detection level. X-ray detection is not a strong function of redshift, X-ray exposure time (at > 1 Ms), power-law slope, or power-law fit probability.

- Almost all (93-98%) of the IRAC-selected power-law galaxies are detected at $24 \mu\text{m}$. The $24 \mu\text{m}$ detection fraction of AGN in the comparison sample increases with X-ray luminosity. Because we require all power-law galaxies to be detected in each of the four IRAC bands, the $24 \mu\text{m}$ detection fraction is almost certainly higher than would be expected for an unbiased sample of AGN. Nevertheless, comparison with a sample suffering from the same bias (the comparison sample) shows that the high detection fraction is representative of an intrinsically X-ray luminous AGN population.
- 30% of the power-law galaxies have radio counterparts from the Richards et al. (2000) VLA survey of the CDF-N. Only two of the 18 radio-detected sources (11%) have radio emission in excess of that predicted by the radio-infrared correlation, suggesting an overall radio-loud fraction of only 3%. In comparison, almost all of the radio-excess AGN of Donley et al. (2005) have optical/MIR SEDs dominated by the stellar bump. Further exploration of a possible systematic difference in the radio properties of the two samples would be interesting. While there is very little overlap between the radio-excess and power-law samples, both have similarly low X-ray detection fractions, suggesting that this may be a common feature of AGN selected independently of their X-ray and optical properties, regardless of redshift or luminosity.
- The optical-MIR SEDs of the power-law galaxies are flatter than the median radio-quiet SED of Elvis et al. (1994). Sources not detected in X-rays have SEDs that drop off more rapidly in the optical than those of the power-law galaxies detected in X-rays, as discussed in Alonso-Herrero et al. (2006), presumably due to increasing optical obscuration. Weakly-detected power-law galaxies have intermediate optical SEDs.
- At least half of the power-law galaxies detected in the X-ray catalog have compact optical counterparts, suggesting that the optical light is dominated by the AGN, as expected. Only 2 non- or weakly-detected sources have compact optical counterparts. 15% of the power-law galaxies in the GOODS field do not have optical counterparts brighter than the GOODS limiting magnitudes.
- Power-law galaxies comprise a subset of the MIR sources selected via color criteria (e.g. Lacy et al. 2004; Stern et al. 2005). While color-selected AGN samples include a higher fraction of high-luminosity AGN than does the power-law selected sample, the color criteria select larger fractions of sources not detected in X-rays, due at least in part to a higher degree of contamination by star-forming galaxies. Combining color-selection with additional tests designed to rule out emission from star-forming galaxies is likely to produce more reliable samples of AGN.
- 68% (2:1) of the X-ray-detected power-law galaxies are obscured ($\log N_{\text{H}}(\text{cm}^{-2}) > 22$), and all of the weakly-detected power-law galaxies are consistent with being obscured, but not Compton-thick ($\log N_{\text{H}}(\text{cm}^{-2}) > 24$). If we assume that all of the X-ray non-detected power-law galaxies are obscured,

we derive a maximum obscured fraction of 81% (4:1). Power-law galaxies also appear to be more heavily obscured at low X-ray luminosities and at high redshift.

- Power-law galaxies detected to high S/N in the IRAC bands account for $\sim 20-30\%$ of both the MIR-detected AGN and the MIR-detected obscured AGN predicted by the X-ray luminosity function synthesis models of Treister et al. (2006) down to $24\text{ }\mu\text{m}$ flux densities of $80\text{ }\mu\text{Jy}$. This percentage is best interpreted as a lower limit since our conservative selection criteria may exclude some power-law objects.
- At all but the highest column densities, NIR/MIR extinction should have only a minor effect on the power-law emission of luminous AGN. IRAC power-law selection, however, is likely to be biased against high-redshift Compton-thick AGN if the covering fractions of the X-ray and NIR-emitting regions are the same.

This work was supported by an NSF Graduate Research Fellowship and by NASA through contract 1255094 issued by JPL/California Institute of Technology. We thank the referee for helpful comments that improved the paper.

APPENDIX

To understand the performance of our power-law AGN selection at high redshift and high luminosity, it is necessary to test it with template spectral energy distributions of appropriately luminous infrared galaxies. It has recently become possible to construct accurate templates in the critical 0.8 to $10\text{ }\mu\text{m}$ range using a combination of Spitzer data, 2MASS total galaxy measurements, and ground-based spectroscopy in the $3-4\text{ }\mu\text{m}$ range. We used these data sources to build templates for 3 ultra-luminous infrared galaxies. Arp 220 and IRAS 17208-0014 are both strongly dominated by star formation, whereas Mrk 273 appears to be powered by a mixture of star formation and an obscured AGN (e.g., Ptak et al. 2003; Farrah et al. 2003).

The references in Table A4 provide total-galaxy flux densities in the optical, near infrared, and at 3.6 and $4.5\text{ }\mu\text{m}$. We derived the near infrared stellar continuum starting from the stellar spectra of Strecker et al. (1979). They have the advantage of being consistently calibrated, are not affected by terrestrial atmospheric absorption, and they extend to $5.5\text{ }\mu\text{m}$. Specifically, we used the spectrum of β And for the near infrared stellar continuum, since its CO absorption approximately matches that of actively star forming galaxies. At wavelengths shortward of $1.2\text{ }\mu\text{m}$, our starting point was the spectral energy distribution of Arp 220 from Silva et al. (1998). For each galaxy, we joined these two spectra at $1.2\text{ }\mu\text{m}$ and then adjusted the normalization and reddening to give a good overall fit through the photometry (UBVR as available plus JHK). The resulting spectrum was taken as the stellar photospheric part of the template (since we need a template valid up to $z = 3$ in the IRAC $3.6\text{ }\mu\text{m}$ band, the behavior at wavelengths shortward of $0.8\text{ }\mu\text{m}$ is unimportant). From Goldader et al. (1995), all three galaxies have strong CO stellar absorption features (bandhead at rest $2.3\text{ }\mu\text{m}$), so the photospheric spectrum is not significantly diluted by other sources at least to $2.4\text{ }\mu\text{m}$. Therefore, we use it to define the template from 0.8 to $2.4\text{ }\mu\text{m}$.

For wavelengths longer than about $6\text{ }\mu\text{m}$, the templates are based on the spectra listed in Table A4. Because the infrared activity is generally concentrated in the nuclei, and the beams used for these spectra are relatively large, we made no corrections for extended emission. To bridge from 2.4 to $6\text{ }\mu\text{m}$, we proceeded as follows. The IRAC photometry at 3.6 and $4.5\text{ }\mu\text{m}$ matches the apertures for the 2MASS total galaxy measurements, so it defines the overall output across the bridge region. The spectrum of M82 (Sturm et al. 2000) indicates that we should expect no strong spectral features between 4 and $6\text{ }\mu\text{m}$ (other than Br α , which should have small equivalent width). To fold in the $3-4\text{ }\mu\text{m}$ region, we used the spectra indicated in Table A4. We estimated the photospheric contribution to these spectra by applying corrections to our stellar template from small aperture photometry in the JHK region, or from small aperture photometry at L, or from both. This contribution was subtracted from the spectrum (using the β And template) and the remainder, representing the excess emission, was added to the total stellar template. This excess spectrum was scaled until the total template matched the photometric point at $3.6\text{ }\mu\text{m}$ (the scaling factors required were small, between 1 and 1.5, indicating that the excess emission was nearly all within the $\sim 1''$ slits used for the spectroscopy). We joined the longest wavelength point from these spectra to the shortest wavelength point in the $6-15\text{ }\mu\text{m}$ spectra with a scaled and adjusted spectrum from M82 (Sturm et al. 2000). We found that adjusting the slope as a power law in wavelength and then normalizing gave a smooth connection that also was compatible with the photometry at $4.5\text{ }\mu\text{m}$. The templates are shown in Figure 20 and are given in Table A5.




FIG. 20.— SED templates for Arp 220 (light solid line), IRAS 17208-0014 (heavy solid line) and Mrk 273 (light dashed line) between 1 and 10 μm . The templates have been normalized to the same value near 7 μm . The influence of the AGN in Mrk 273 is apparent both in the reduced equivalent widths of the aromatic features and in the filling-in of the SED near 5 μm .

REFERENCES

- Akiyama, M., Ueda, Y., Ohta, K., Takahashi, T., & Yamada, T. 2003, *ApJS*, 148, 275
- Akylas, A., Georgantopoulos, I., Georgakakis, A., Kitsionas, S., & Hatziminaoglou, E., 2006, *A&A*, in press
- Alexander, D. M., Aussel, H., Bauer, F. E., Brandt, W. N., Hornschemeier, A. E., Vignali, C., Garmire, G. P., & Schneider, D. P. 2002, *ApJ*, 568, L85
- Alexander, D. M., et al. 2003, *AJ*, 126, 539
- Alexander, D. M., Smail, I., Bauer, F. E., Chapman, S. C., Blain, A. W., Brandt, W. N., & Ivison, R. J. 2005, *Nature*, 434, 738
- Alonso-Herrero, A., et al. 2004, *ApJS*, 154, 155
- Alonso-Herrero, A., et al. 2006, *ApJ*, 640, 167
- Appleton, P. N., et al. 2004, *ApJS*, 154, 147
- Armus, L. et al. 2006, submitted to *ApJ*, and Armus, private communication
- Ballantyne, D. R., Everett, J. E., & Murray, N. 2006, *ApJ*, 639, 740
- Barger, A. J., Cowie, L. L., Bautz, M. W., Brandt, W. N., Garmire, G. P., Hornschemeier, A. E., Ivison, R. J., & Owen, F. N. 2001, *AJ*, 122, 2177
- Barger, A. J., Cowie, L. L., Brandt, W. N., Capak, P., Garmire, G. P., Hornschemeier, A. E., Steffen, A. T., & Wehner, E. H. 2002, *AJ*, 124, 1839
- Barger, A. J., et al. 2003, *AJ*, 126, 632
- Barger, A. J., Cowie, L. L., Mushotzky, R. F., Yang, Y., Wang, W.-H., Steffen, A. T., & Capak, P. 2005, *AJ*, 129, 578
- Barmby, P., et al. 2006, *ApJ*, 642, 126
- Bauer, F. E., Alexander, D. M., Brandt, W. N., Schneider, D. P., Treister, E., Hornschemeier, A. E., & Garmire, G. P. 2004, *AJ*, 128, 2048
- Bell, E. F. 2003, *ApJ*, 586, 794
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bevington, P. R., & Robinson, D. K. 2003, *Data reduction and error analysis for the physical sciences*, 3rd ed., by Philip R. Bevington, and Keith D. Robinson. Boston, MA: McGraw-Hill, ISBN 0-07-247227-8, 2003.
- Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, *ApJ*, 224, 132
- Brand, K., et al. 2006, *ApJ*, 644, 143
- Brusa, M., et al. 2003, *A&A*, 409, 65
- Capak, P., et al. 2004, *AJ*, 127, 180
- Chapman, S. C., Blain, A. W., Smail, I., & Ivison, R. J. 2005, *ApJ*, 622, 772
- Comastri, A., Fiore, F., Vignali, C., Matt, G., Perola, G. C., & La Franca, F. 2001, *MNRAS*, 327, 781
- Condon, J. J., Cotton, W. D., & Broderick, J. J. 2002, *AJ*, 124, 675
- Cowie, L. L., Barger, A. J., Hu, E. M., Capak, P., & Songaila, A. 2004, *AJ*, 127, 3137
- Dale, D. A., & Helou, G. 2002, *ApJ*, 576, 159
- Dawson, S., McCrady, N., Stern, D., Eckart, M. E., Spinrad, H., Liu, M. C., & Graham, J. R. 2003, *AJ*, 125, 1236
- Devriendt, J. E. G., Guiderdoni, B., & Sadat, R. 1999, *A&A*, 350, 381
- Donley, J. L., Rieke, G. H., Rigby, J. R., & Pérez-González, P. G. 2005, *ApJ*, 634, 169
- Duc, P.-A., Mirabel, I. F., and Maza, J. 1997, *A&AS*, 124, 533
- Elvis, M., et al. 1994, *ApJS*, 95, 1
- Fazio, G. G., et al. 2004, *ApJS*, 154, 10
- Farrah, D., Alfonso, J., Efstathiou, A., Rowan-Robinson, M., Fox, M., & Clements, D. 2003, *MNRAS*, 343, 585
- Franceschini, A., et al. 2005, *AJ*, 129, 2074
- Freeman, P. E., Kashyap, V., Rosner, R., & Lamb, D. Q. 2002, *ApJS*, 138, 185
- Frueh, M. L., Corwin, H. G. Jr., de Vaucouleurs, G., & Buta, R. 1996, *AJ*, 111, 722
- Giacconi, R., et al. 2001, *ApJ*, 551, 624
- Giacconi, R., et al. 2002, *ApJS*, 139, 369
- Giavalisco, M., et al. 2004, *ApJ*, 600, L93
- Gilli, R., Salvati, M., & Hasinger, G. 2001, *A&A*, 366, 407
- Gilli, R., et al. 2003, *ApJ*, 592, 721
- Gilli, R. 2004, *Advances in Space Research*, 34, 2470
- Goldader, J. D., Joseph, R. D., Doyon, Rene, & Sanders, D. B. 1995, *ApJ*, 444, 97
- Gordon, K. D., et al. 2004, *Proc. SPIE*, 5487, 177
- Gordon, K. D., et al. 2005, *PASP*, 117, 503
- Granato, G. L., Danese, L., & Franceschini, A. 1997, *ApJ*, 486, 147
- Hasinger, G. 2004, *Nuclear Physics B Proceedings Supplements*, 132, 86
- Hatziminaoglou, E., et al. 2005, *AJ*, 129, 1198
- Higdon, S. J. U., Armus, L., Higdon, J. L., Soifer, B. T., & Spoon, H. W. W., 2006, *ApJ*, in press, astro-ph/0605359
- Hornschemeier, A. E., et al. 2001, *ApJ*, 554, 742
- Hornschemeier, A. E., et al. 2003, *AJ*, 126, 575
- Houck, J. R., et al. 2005, *ApJ*, 622, L105
- Huang, J.-S., et al. 2004, *ApJS*, 154, 44
- Imanishi, M. 2006, *AJ*, 131, 2406
- Imanishi, M., Dudley, C. C., & Maloney, P. R. 2006, *ApJ*, 637, 114
- Ivezić, Ž., et al. 2002, *AJ*, 124, 2364
- Ivison, R. J., et al. 2004, *ApJS*, 154, 124
- Johansson, P. H., Väisänen, P., & Vaccari, M. 2004, *A&A*, 427, 795
- Kellermann, K. I., Sramek, R., Schmidt, M., Shaffer, D. B., & Green, R. 1989, *AJ*, 98, 1195

- Klaas, U., Haas, M., Heinrichsen, I., & Schulz, B. 1997, *Å*, 325, L21
- Koekemoer, A. M., et al. 2004, *ApJ*, 600, L123
- Kuraszkiewicz, J. et al. 2007, *ApJ*, submitted
- La Franca, F., et al. 2005, *ApJ*, 635, 864
- Lacy, M., et al. 2004, *ApJS*, 154, 166
- Lacy, M., Canalizo, G., Rawlings, S., Sajina, A., Storrie-Lombardi, L., Armus, L., Marleau, F. R., & Muzzin, A. 2005, *Memorie della Societa Astronomica Italiana*, 76, 154
- Lawrence, A. 1991, *MNRAS*, 252, 586
- Leipski, C., et al. 2005, *A&A*, 440, L5
- Maccacaro, T., Perola, G. C., & Elvis, M. 1982, *ApJ*, 257, 47
- Maccacaro, T., Gioia, I. M., Wolter, A., Zamorani, G., & Stocke, J. T. 1988, *ApJ*, 326, 680
- Maiolino, R., & Rieke, G. H. 1995, *ApJ*, 454, 95
- Maiolino, R., Salvati, M., Bassani, L., Dadina, M., della Ceca, R., Matt, G., Risaliti, G., & Zamorani, G. 1998, *A&A*, 338, 781
- Maiolino, R., Marconi, A., Salvati, M., Risaliti, G., Severgnini, P., Oliva, E., La Franca, F., & Vanzi, L. 2001a, *A&A*, 365, 28
- Maiolino, R., Marconi, A., & Oliva, E. 2001b, *A&A*, 365, 37
- Marble, A. R., Hines, D. C., Schmidt, G. D., Smith, P. S., Surace, J. A., Armus, L., Cutri, R. M., & Nelson, B. O. 2003, *ApJ*, 590, 707
- Martínez-Sansigre, A., Rawlings, S., Lacy, M., Fadda, D., Marleau, F. R., Simpson, C., Willott, C. J., & Jarvis, M. J. 2005, *Nature*, 436, 666
- Martínez-Sansigre, A., Rawlings, S., Lacy, M., Fadda, D., Jarvis, M. J., Marleau, F. R., Simpson, C., Willott, C. J. 2006, *MNRAS*, in press
- McLeod, K. K., & Rieke, G. H. 1994, *ApJ*, 431, 137
- Morrison, G., et al. 2006, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0603412
- Mushotzky, R. F., Cowie, L. L., Barger, A. J., & Arnaud, K. A. 2000, *Nature*, 404, 459
- Neugebauer, G., Oke, J. B., Becklin, E. E., & Matthews, K. 1979, *ApJ*, 230, 79
- Origlia, L., Moorwood, A. F. M., & Oliva, E. 1993, *A&A*, 280, 536
- Page, M. J., et al. 2003, *Astronomische Nachrichten*, 324, 101
- Papovich, C., et al. 2004, *ApJS*, 154, 70
- Pearson, C. 2005, *MNRAS*, 358, 1417
- Percival, W. J., Miller, L., McLure, R. J., & Dunlop, J. S. 2001, *MNRAS*, 322, 843
- Pérez-González, P. G., et al. 2005, *ApJ*, 630, 82
- Perola, G. C., et al. 2004, *A&A*, 421, 491
- Polletta, M. d. C., et al. 2006, *ApJ*, 642, 673
- Ptak, A., Heckman, T., Levenson, N. A., Weaver, K., & Strickland, D. 2003, *ApJ*, 592, 782
- Reichert, G. A., Mushotzky, R. F., Holt, S. S., & Petre, R. 1985, *ApJ*, 296, 69
- Rieke, G. H. 1978, *ApJ*, 226, 550
- Rieke, G. H., & Lebofsky, M. J. 1981, *ApJ*, 250, 87
- Rieke, G. H., & Lebofsky, M. J. 1985, *ApJ*, 288, 618
- Rieke, G. H., Cutri, R. M., Black, J. H., Kailey, W. F., McAlary, C. W., Lebofsky, M. J., & Elston, R. 1985, *ApJ*, 290, 116
- Richards, E. A. 2000, *ApJ*, 533, 611
- Richards, G. T., et al. 2006, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0601558
- Rigby, J. R., et al. 2004, *ApJS*, 154, 160
- Rigby, J. R., Rieke, G. H., Pérez-González, P. G., Donley, J. L., Alonso-Herrero, A., Huang, J.-S., Barmby, P., & Fazio, G. G. 2005, *ApJ*, 627, 134
- Rigopoulou, D., Spoon, H. W. W., Genzel, R., Lutz, D., Moorwood, A. F. M., & Tran, Q. D. 1999, *AJ*, 118, 2625
- Risaliti, G., Maiolino, R., & Salvati, M. 1999, *ApJ*, 522, 157
- Risaliti, G., et al. 2006, *MNRAS*, 365, 303
- Sajina, A., Lacy, M., & Scott, D. 2005, *ApJ*, 621, 256
- Scoville, N. Z., Evans, A. S., Thompson, R., Rieke, M., Hines, D. C., Low, F. J., Dinshaw, N., Surace, J. A., and Armus, L. 2000, *AJ*, 119, 991
- Shi, Y. et al. 2006, *ApJ*, submitted
- Silva, L., Granato, G. L., Bressan, A., & Danese, L. 1998, *ApJ*, 509, 103
- Spinoglio, L., & Malkan, M. A. 1989, *ApJ*, 342, 83
- Steffen, A. T., Barger, A. J., Cowie, L. L., Mushotzky, R. F., & Yang, Y. 2003, *ApJ*, 596, L23
- Stern, D., Djorgovski, S. G., Perley, R. A., de Carvalho, R. R., & Wall, J. V. 2000, *AJ*, 119, 1526
- Stern, D., et al. 2005, *ApJ*, 631, 163
- Strecker, D. W., Erickson, E. F., & Witteborn, F. C. 1979, *ApJS*, 41, 501
- Sturm, E., Lutz, D., Tran, D., Feuchtgruber, H., Genzel, R., Kunze, D., Moorwood, A. F. M., & Thornley, M. D. 2000, *A&A*, 358, 481
- Surace, J. A., Sanders, D. B., & Evans, A. E. 2000, *ApJ*, 529, 170
- Swinbank, A. M., Smail, I., Chapman, S. C., Blain, A. W., Ivison, R. J., & Keel, W. C. 2004, *ApJ*, 617, 64
- Szokoly, G. P., et al. 2004, *ApJS*, 155, 271
- Tozzi, P., et al. 2006, *A&A*, 451, 457
- Treister, E., et al. 2004, *ApJ*, 616, 123
- Treister, E., et al. 2005, *ApJ*, 621, 104
- Treister, E., & Urry, C. M. 2005, *ApJ*, 630, 115
- Ueda, Y., Akiyama, M., Ohta, K., & Miyaji, T. 2003, *ApJ*, 598, 886
- Veilleux, S., Kim, D.-C., & Sanders, D. B. 1999, *ApJ*, 522, 113
- Weedman, D. W., Le Floc'h, E., Higdon, S. J. U., Higdon, J. L., & Houck, J. R. 2006, *ApJ*, 638, 613
- Wilkes, B. J., Pounds, K. A., Schmidt, G. D., Smith, P. S., Cutri, R. M., Ghosh, H., Nelson, B., & Hines, D. C. 2005, *ApJ*, 634, 183
- Wirth, G. D., et al. 2004, *AJ*, 127, 3121
- Worsley, M. A., Fabian, A. C., Barcons, X., Mateos, S., Hasinger, G., & Brunner, H. 2004, *MNRAS*, 352, L28
- Worsley, M. A., et al. 2005, *MNRAS*, 357, 1281
- Worsley, M. A., Fabian, A. C., Bauer, F. E., Alexander, D. M., Brandt, W. N., & Lehmer, B. D. 2006, *MNRAS*, 368, 1735
- Yan, L., et al. 2004, *ApJS*, 154, 60
- Yan, L., et al. 2005, *ApJ*, 628, 604
- Yun, M. S., Reddy, N. A., & Condon, J. J. 2001, *ApJ*, 554, 803

TABLE 1
POWER-LAW GALAXY SAMPLE

ID	α_{2000}	δ_{2000}	P_X^a	α^b	z	24 μ m Flux Density (μ Jy)	1.4 GHz Flux Density (μ Jy)	0.5-8 keV X-ray Flux (ergs s $^{-1}$ cm $^{-2}$)	log L $_X^c$ (ergs s $^{-1}$)	X-ray Exposure (s)	θ^d (')	q^e
CDFN 11348	12 35 25.58	62 16 09.41	0.86	-0.86	...	204.0	< 80.0	< 8.10E-16	...	9.59E+05	9.6	> 0.41
CDFN 17315	12 35 29.37	62 12 56.53	0.22	-1.94	2.41	251.9	135.0	7.02E-15	44.5	7.43E+05	8.9	0.27
CDFN 11516	12 35 37.11	62 17 23.40	0.49	-0.89	2.05	234.9	< 80.0	5.27E-15	44.2	9.00E+05	8.7	> 0.47
CDFN 07514	12 35 38.08	62 19 40.56	0.86	-0.79	2.97 $\pm$ 0.40	< 157.2	< 80.0	2.00E-15	44.2	9.86E+05	9.7	...
CDFN 12939	12 35 38.52	62 16 42.91	0.69	-1.15	0.71	1413.8	3550.0	2.23E-13	44.7	9.60E+05	8.3	-0.40
CDFN 14642	12 35 43.35	62 16 17.15	0.84	-1.17	2.36 $\pm$ 0.34	< 117.6	< 80.0	9.19E-16	43.6	1.25E+06	7.6	...
CDFN 08362	12 35 46.64	62 20 13.23	0.57	-1.39	...	118.5	< 80.0	< 7.85E-16	...	9.66E+05	9.3	> 0.17
CDFN 10934	12 35 48.89	62 19 04.46	0.26	-1.58	...	115.4	< 80.0	< 4.92E-16	...	9.84E+05	8.4	> 0.16
CDFN 16796	12 35 49.43	62 15 36.52	0.22	-0.79	2.20	585.6	74.6	1.34E-15	43.7	1.58E+06	6.8	0.89
CDFN 25218	12 35 50.36	62 10 41.70	0.11	-2.04	...	< 108.5	< 80.0	4.21E-16	...	7.04E+05	7.2	...
CDFN 20981	12 35 54.30	62 13 43.37	0.95	-0.69	...	131.7	< 80.0	< 2.04E-16	...	1.74E+06	6.0	> 0.22
CDFN 25248	12 36 06.37	62 12 32.41	0.81	-0.60	2.89 $\pm$ 0.39	139.2	< 80.0	9.64E-16	43.8	1.81E+06	4.8	> 0.24
CDFN 33052	12 36 08.87	62 08 03.56	0.10	-1.37	...	151.8	< 80.0	< 4.67E-16	...	5.43E+05	7.3	> 0.28
CDFN 27463	12 36 11.95	62 11 47.18	0.77	-0.53	2.92 $\pm$ 0.39	180.3	< 80.0	< 1.23E-16	< 42.9	1.68E+06	4.5	> 0.35
CDFN 31054	12 36 14.04	62 09 47.93	0.78	-0.59	1.65 $\pm$ 0.27	60.8	< 80.0	< 2.46E-16	< 42.6	1.79E+06	5.6	> -0.12
CDFN 23661	12 36 22.98	62 15 26.28	0.16	-1.40	2.59	454.5	< 80.0	1.88E-14	45.0	1.85E+06	3.0	> 0.75
CDFN 39358	12 36 23.37	62 06 05.23	0.21	-1.92	1.86 $\pm$ 0.29	479.5	71.7	4.71E-16	43.0	5.42E+05	8.3	0.83
CDFN 37928	12 36 29.21	62 07 37.61	0.91	-0.59	1.35 $\pm$ 0.23	161.6	< 80.0	< 2.15E-16	< 42.3	1.77E+06	6.6	> 0.31
CDFN 37999	12 36 32.59	62 07 58.95	0.11	-2.14	2.00	711.8	90.6	1.83E-15	43.7	1.78E+06	6.2	0.90
CDFN 27641	12 36 35.60	62 14 23.51	0.86	-1.82	2.02	1345.2	87.8	2.52E-15	43.8	1.92E+06	1.2	1.19
CDFN 28773	12 36 36.66	62 13 46.46	0.63	-0.74	0.96	459.6	< 80.0	1.01E-14	43.7	1.91E+06	1.1	> 0.76
CDFN 14667	12 36 36.90	62 22 27.27	0.57	-2.03	...	292.9	< 80.0	3.45E-15	...	1.06E+06	8.6	> 0.56
CDFN 37334	12 36 37.08	62 08 51.90	0.32	-0.90	1.71 $\pm$ 0.27	301.7	71.1	1.19E-16	42.3	1.78E+06	5.2	0.63
CDFN 44902	12 36 40.59	62 04 51.34	0.27	-0.87	1.16 $\pm$ 0.22	166.6	< 80.0	< 8.28E-16	< 42.8	6.96E+05	9.1	> 0.32
CDFN 24409	12 36 42.22	62 17 10.89	0.65	-0.80	2.72	91.7	< 80.0	3.35E-15	44.3	1.65E+06	3.2	> 0.06
CDFN 11965	12 36 42.23	62 24 39.01	0.59	-0.71	1.90 $\pm$ 0.29	480.4	82.1	1.95E-15	43.7	9.62E+05	10.7	0.77
CDFN 14360	12 36 47.27	62 23 49.06	0.29	-1.28	...	216.0	< 80.0	< 7.05E-16	...	9.75E+05	9.9	> 0.43
CDFN 18906	12 36 48.30	62 21 06.89	0.76	-1.45	...	149.4	< 80.0	< 2.94E-16	...	1.68E+06	7.2	> 0.27
CDFN 41981	12 36 49.66	62 07 37.84	0.50	-2.30	1.54 $\pm$ 0.25	1239.3	307.0	2.59E-14	44.6	1.74E+06	6.3	0.61
CDFN 49937	12 36 58.76	62 04 01.80	0.12	-1.17	0.29	138.7	< 80.0	9.75E-15	42.4	7.33E+05	10.1	> 0.24
CDFN 19080	12 36 58.96	62 22 14.89	0.28	-1.94	...	106.6	< 80.0	< 4.41E-16	...	1.62E+06	8.4	> 0.12
CDFN 14046	12 36 59.07	62 25 21.81	0.31	-1.04	2.82 $\pm$ 0.38	164.3	< 80.0	1.16E-14	44.9	8.68E+05	11.5	> 0.31
CDFN 44836	12 37 01.69	62 07 20.21	0.56	-3.15	...	323.6	< 80.0	8.56E-16	...	1.74E+06	6.9	> 0.61
CDFN 53291	12 37 01.70	62 02 22.05	0.10	-1.59	2.17 $\pm$ 0.32	99.6	< 80.0	< 1.89E-15	< 43.8	6.41E+05	11.8	> 0.10
CDFN 45610	12 37 06.59	62 07 26.83	0.79	-1.05	...	-1.0	< 80.0	< 7.08E-16	...	1.76E+06	7.0	...
CDFN 29120	12 37 06.90	62 17 02.37	0.29	-1.06	1.02	766.3	< 80.0	3.54E-14	44.3	1.87E+06	3.9	> 0.98
CDFN 16936	12 37 07.01	62 24 27.81	0.92	-1.25	...	161.3	< 80.0	< 1.00E-15	...	1.08E+06	10.8	> 0.30
CDFN 45284	12 37 09.85	62 08 00.82	0.39	-1.40	2.18	128.6	< 80.0	2.11E-14	44.9	1.71E+06	6.6	> 0.21
CDFN 20246	12 37 10.00	62 22 58.92	0.20	-1.42	1.41 $\pm$ 0.24	318.6	708.0	5.21E-15	43.8	1.60E+06	9.4	-0.35
CDFN 38580	12 37 12.07	62 12 11.61	0.70	-1.14	2.91	< 80.0	< 80.0	3.84E-16	43.4	1.72E+06	3.5	...
CDFN 32673	12 37 13.72	62 15 45.14	0.52	-1.14	...	129.8	< 80.0	4.46E-16	...	1.90E+06	3.7	> 0.21
CDFN 43967	12 37 14.06	62 09 16.80	0.38	-0.96	1.84 $\pm$ 0.28	105.3	< 80.0	3.57E-15	43.9	1.83E+06	5.7	> 0.12
CDFN 30147	12 37 16.69	62 17 33.18	0.56	-1.75	1.15	1015.7	346.0	2.17E-14	44.2	1.84E+06	5.1	0.47
CDFN 50281	12 37 16.99	62 05 53.05	0.59	-1.05	1.94	398.5	< 80.0	1.94E-15	43.7	7.26E+05	8.9	> 0.70
CDFN 28149	12 37 17.90	62 18 55.56	0.17	-1.24	2.24	195.2	< 80.0	9.95E-15	44.6	1.78E+06	6.2	> 0.39
CDFN 46227	12 37 21.72	62 08 50.41	0.84	-0.80	2.56 $\pm$ 0.36	292.1	< 80.0	< 2.43E-16	< 43.1	1.77E+06	6.6	> 0.56
CDFN 51788	12 37 23.03	62 05 39.45	0.99	-1.08	2.28 $\pm$ 0.33	146.6	78.5	< 2.65E-15	< 44.0	7.43E+05	9.4	0.27
CDFN 27360	12 37 26.58	62 20 26.46	0.14	-2.26	1.86 $\pm$ 0.29	777.7	102.0	1.40E-15	43.5	1.40E+06	8.0	0.88
CDFN 38126	12 37 28.65	62 14 22.56	0.58	-1.80	...	226.1	< 80.0	1.20E-16	...	1.82E+06	5.0	> 0.45
CDFN 40913	12 37 30.77	62 12 58.50	0.97	-1.78	...	139.3	107.0	< 2.08E-16	...	1.79E+06	5.3	0.11

TABLE 1
— CONTINUED

ID	α_{2000}	δ_{2000}	P_{χ} ^a	α ^b	z	24 μm Flux Density (μJy)	1.4 GHz Flux Density (μJy)	0.5-8 keV X-ray Flux ($\text{ergs s}^{-1} \text{ cm}^{-2}$)	$\log L_x$ ^c (ergs s^{-1})	X-ray Exposure (s)	θ ^d (')	q ^e
CDFN 39529	12 37 36.83	62 14 28.72	0.20	-1.44	...	118.9	57.8	6.21E-15	...	1.74E+06	6.0	0.31
CDFN 26243	12 37 39.78	62 22 39.67	0.38	-2.08	...	218.2	< 80.0	< 3.33E-15	...	6.12E+05	10.7	> 0.44
CDFN 44598	12 37 41.00	62 12 00.29	0.26	-0.63	1.17	558.2	< 80.0	2.02E-14	44.2	1.76E+06	6.7	> 0.84
CDFN 34143	12 37 42.58	62 18 11.67	0.16	-1.55	2.31	149.6	< 80.0	2.08E-14	44.9	1.69E+06	7.8	> 0.27
CDFN 44751	12 37 44.67	62 12 18.71	0.21	-0.94	1.69 ± 0.27	416.1	67.3	2.88E-16	42.7	1.76E+06	7.1	0.79
CDFN 40112	12 37 57.32	62 16 27.57	0.28	-1.55	2.92	122.3	< 80.0	4.68E-15	44.5	1.70E+06	8.7	> 0.18
CDFN 54722	12 37 57.55	62 08 00.51	0.65	-1.13	...	174.9	< 80.0	< 1.01E-15	...	6.41E+05	10.3	> 0.34
CDFN 49940	12 37 59.62	62 11 02.09	0.58	-1.42	0.91	2132.0	85.0	1.52E-13	44.8	7.33E+05	9.1	1.40
CDFN 45792	12 38 00.94	62 13 35.90	0.43	-0.72	0.44	4092.5	190.0	7.78E-14	43.7	1.44E+06	8.8	1.33
CDFN 53792	12 38 31.18	62 12 21.98	0.59	-0.77	1.63 ± 0.26	297.0	< 80.0	3.76E-15	43.8	6.62E+05	12.4	> 0.57
CDFN 53753	12 38 32.19	62 12 30.60	0.25	-0.66	1.63 ± 0.26	198.4	< 80.0	< 2.79E-15	< 43.7	6.15E+05	12.5	> 0.39
CDFN 54888	12 38 34.09	62 12 05.88	0.67	-1.75	1.96 ± 0.30	476.4	< 80.0	< 1.59E-15	< 43.6	5.57E+05	12.8	> 0.77

^a Chi-squared probability (see §3) ^b IRAC spectra index; $f_{\nu} \propto \nu^{\alpha}$ ^c 0.5-8 keV ^d X-ray off-axis angle ^e $q = \log(f_{24 \mu\text{m}}/f_{1.4 \text{ GHz}})$

TABLE 2
X-RAY PROPERTIES OF X-RAY WEAKLY-DETECTED POWER-LAW GALAXIES

Source	z	X-ray Band ^a	EER (%)	Detection σ	Aperture Corrected Count Rate (s ⁻¹)	Γ	Flux (erg s ⁻¹ cm ⁻²)	H/S ^b	log N _H (cm ⁻²)
CDFN14642	2.36	Full	80	4.6	4.44E-05	0.28	9.19E-16	11.17	23.7 ^{+0.1} _{-0.1}
		Hard	70	3.3	2.33E-05		6.98E-16		
		Soft	60	3.3	1.27E-05		6.08E-17		
CDFN20981	...	Full	80	≤ 2.5	$\leq 1.44E-05$	≥ 0.41	$\leq 1.68E-16$	≤ 6.96	$\leq 22.5^{+0.1}_{-0.1} - \leq 23.7^{+0.1}_{-0.1}$
		Hard	80	≤ 2.5	$\leq 1.00E-05$		$\leq 2.31E-16$		
		Soft	60	3.3	6.35E-06		3.18E-17		
CDFN25218	...	Full	80	3.5	3.61E-05	≤ 0.61	4.21E-16	≥ 5.58	$\geq 22.4^{+0.1}_{-0.1} - \geq 23.6^{+0.1}_{-0.1}$
		Hard	70	2.8	2.35E-05		5.42E-16		
		Soft	80	≤ 2.5	$\leq 1.86E-05$		$\leq 9.33E-17$		
CDFN37334	1.71	Full	80	2.6	1.02E-05	≥ 0.20	1.19E-16	≤ 8.83	$\leq 23.3^{+0.1}_{-0.1}$
		Hard	80	≤ 2.5	$\leq 1.17E-05$		$\leq 2.70E-16$		
		Soft	70	3.2	5.85E-06		2.93E-17		
CDFN37928	1.35	Full	70	≤ 2.5	$\leq 9.22E-06$	≥ 0.69	$\leq 1.07E-16$	≤ 5.09	$\leq 22.9^{+0.1}_{-0.1}$
		Hard	70	≤ 2.5	$\leq 8.10E-06$		$\leq 1.87E-16$		
		Soft	60	2.8	7.03E-06		3.52E-17		
CDFN38126	...	Full	80	2.7	1.03E-05	...	1.20E-16	...	...
		Hard	80	≤ 2.5	$\leq 1.03E-05$		$\leq 2.38E-16$		
		Soft	80	≤ 2.5	$\leq 6.97E-06$		$\leq 3.49E-17$		
CDFN39358	1.86	Full	80	2.8	4.04E-05	≤ 0.14	4.71E-16	≥ 9.42	$\geq 23.4^{+0.1}_{-0.1}$
		Hard	80	2.6	3.39E-05		7.82E-16		
		Soft	70	≤ 2.5	$\leq 1.59E-05$		$\leq 7.97E-17$		
CDFN44751	1.69	Full	80	3.7	2.47E-05	≥ 0.52	2.88E-16	≤ 6.21	$\leq 23.2^{+0.1}_{-0.1}$
		Hard	80	≤ 2.5	$\leq 2.01E-05$		$\leq 4.64E-16$		
		Soft	80	4.2	1.43E-05		7.17E-17		
CDFN44902	1.16	Full	70	≤ 2.5	$\leq 4.54E-05$	≥ 1.05	$\leq 5.29E-16$	≤ 3.41	$\leq 22.5^{+0.1}_{-0.2}$
		Hard	80	≤ 2.5	$\leq 2.78E-05$		$\leq 6.41E-16$		
		Soft	60	5.0	3.60E-05		1.80E-16		
CDFN46227	2.56	Full	60	≤ 2.5	$\leq 1.04E-05$	≥ 0.83	$\leq 1.21E-16$	≤ 4.37	$\leq 23.3^{+0.1}_{-0.2}$
		Hard	60	≤ 2.5	$\leq 6.83E-06$		$\leq 1.58E-16$		
		Soft	60	2.7	6.91E-06		3.46E-17		

^a Full band = 0.5-8 keV; Hard band = 2-8 keV; Soft band = 0.5-2 keV ^b Flux ratio corrected for Galactic absorption

TABLE 3
X-RAY PROPERTIES OF X-RAY-DETECTED POWER-LAW AGN

Source	z	0.5-8 keV Flux ^a (erg s ⁻¹ cm ⁻²)	2-8 keV Flux ^a (erg s ⁻¹ cm ⁻²)	0.5-2 keV Flux ^a (erg s ⁻¹ cm ⁻²)	Γ^a	H/S ^b	log N _H ^c (cm ⁻²)
CDFN07514	2.97	2.00E-15	1.54E-15	4.19E-16	1.1	3.54	23.3 ^{+0.1} _{-0.2}
CDFN11516	2.05	5.27E-15	2.85E-15	2.04E-15	1.8	1.33	21.2 ^{+1.0} _{...}
CDFN11965	1.90	1.95E-15	1.56E-15	3.44E-16	0.9	4.38	23.1 ^{+0.1} _{-0.2}
CDFN12939	0.71	2.23E-13	1.36E-13	8.05E-14	1.6	1.62	21.4 ^{+0.4} _{...}
CDFN14046	2.82	1.16E-14	6.63E-15	4.32E-15	1.7	1.47	22.0 ^{+0.6} _{...}
CDFN14667	...	3.45E-15	2.13E-15	1.13E-15	1.6	1.81	21.5 ^{+0.3} _{-1.3} - 22.6 ^{+0.3} _{-0.9}
CDFN16796	2.20	1.34E-15	1.20E-15	1.20E-16	0.4	9.72	23.6 ^{+0.1} _{-0.1}
CDFN17315	2.41	7.02E-15	4.72E-15	2.18E-15	1.5	2.08	22.6 ^{+0.3} _{-0.4}
CDFN20246	1.41	5.21E-15	4.99E-15	1.35E-16	-0.6	36.19	23.6 ^{+0.0} _{-0.0}
CDFN23661	2.59	1.88E-14	1.10E-14	7.43E-15	1.8	1.41	21.8 ^{+0.7} _{...}
CDFN24409	2.72	3.35E-15	2.32E-15	9.48E-16	1.4	2.35	22.8 ^{+0.2} _{-0.3}
CDFN25248	2.89	9.64E-16	8.63E-16	1.08E-16	0.5	7.75	23.7 ^{+0.1} _{-0.1}
CDFN27360	1.86	1.40E-15	1.32E-15	1.52E-16	0.5	8.43	23.4 ^{+0.1} _{-0.1}
CDFN27641	2.02	2.52E-15	2.48E-15	2.07E-16	0.2	11.66	23.6 ^{+0.1} _{-0.1}
CDFN28149	2.24	9.95E-15	6.20E-15	3.43E-15	1.6	1.73	22.3 ^{+0.3} _{-1.2}
CDFN28773	0.96	1.01E-14	5.35E-15	4.57E-15	1.9	1.12	≤ 21.2
CDFN29120	1.02	3.54E-14	1.89E-14	1.57E-14	1.9	1.15	≤ 21.4
CDFN30147	1.15	2.17E-14	1.73E-14	4.26E-15	1.0	3.92	22.6 ^{+0.1} _{-0.2}
CDFN32673	...	4.46E-16	5.30E-16	≤ 2.55E-17	≤ 1.4	≥ 19.96	≥ 22.9 ^{+0.0} _{-0.1} - ≥ 24.0 ^{+0.0} _{-0.1}
CDFN34143	2.31	2.08E-14	1.29E-14	7.30E-15	1.6	1.69	22.2 ^{+0.4} _{...}
CDFN37999	2.00	1.83E-15	1.83E-15	1.17E-16	0.1	15.24	23.7 ^{+0.1} _{-0.1}
CDFN38580	2.91	3.84E-16	3.74E-16	2.93E-17	0.2	12.42	23.9 ^{+0.1} _{-0.1}
CDFN39529	...	6.21E-15	5.16E-15	1.09E-15	0.9	4.57	22.3 ^{+0.1} _{-0.1} - 23.5 ^{+0.1} _{-0.1}
CDFN40112	2.92	4.68E-15	2.42E-15	2.27E-15	2.0	1.02	≤ 20.7
CDFN41981	1.54	2.59E-14	2.18E-14	4.34E-15	0.9	4.85	23.0 ^{+0.1} _{-0.1}
CDFN43967	1.84	3.57E-15	2.44E-15	1.02E-15	1.4	2.30	22.5 ^{+0.2} _{-0.3}
CDFN44598	1.17	2.02E-14	1.28E-14	6.96E-15	1.6	1.76	21.8 ^{+0.3} _{-2.2}
CDFN44836	...	8.56E-16	9.77E-16	≤ 5.71E-17	≤ 1.4	≥ 16.43	≥ 22.8 ^{+0.1} _{-0.1} - ≥ 24.0 ^{+0.1} _{-0.1}
CDFN45284	2.18	2.11E-14	1.27E-14	7.82E-15	1.7	1.55	22.1 ^{+0.4} _{...}
CDFN45792	0.44	7.78E-14	3.83E-14	3.87E-14	2.0	0.94	...
CDFN49937	0.29	9.75E-15	5.69E-15	3.65E-15	1.7	1.49	21.0 ^{+0.4} _{...}
CDFN49940	0.91	1.52E-13	9.11E-14	5.64E-14	1.7	1.54	21.4 ^{+0.5} _{...}
CDFN50281	1.94	1.94E-15	9.82E-16	7.57E-16	1.8	1.24	≤ 22.0
CDFN53792	1.63	3.76E-15	3.06E-15	6.42E-16	0.9	4.60	23.0 ^{+0.1} _{-0.1}

^a Alexander et al. (2003) ^b Flux ratio corrected for Galactic absorption ^c Sources with no lower error have an immeasurably low lower limit.

TABLE 4
A4: INPUT DATA FOR TEMPLATES

Galaxy	UBVR	JHK	3–4 μ m spectrum	3.6, 4.5 μ m	6–15 μ m spectrum	Other
Arp 220	1	4	5	7; 54.2, 46.6 mJy	9	12, 13
IRAS 17208-0014	2	4	6	8; 17.9, 17.3 mJy	10	14
Mrk 273	3	4	5	7; 27.6, 38.2 mJy	11	15

REFERENCES. — (1) Frueh et al. 1996; (2) Duc, Mirabel, & Maza 1997; (3) Surace, Sanders, & Evans 2000; (4) 2MASS via NED; (5) Imanishi, Dudley, & Maloney 2006; (6) Imanishi 2006; (7) Spitzer PID 32, extraction aperture diameter of 90" for Arp 220 and 50" for Mrk 273; (8) Spitzer PID 3672, extraction aperture 30" in diameter; (9) Armus et al. 2006; (10) Rigopoulou et al. 1999; (11) Higdon et al. 2006; (12) Klaas et al. 1997; (13) Rieke et al. 1985; (14) Scoville et al. 2000; (15) Rieke 1978

TABLE 5
A5: TEMPLATES

Arp 220		IRAS 17208-0014		Mrk 273	
$\lambda(\mu\text{m})$	$f_\nu(\text{Jy})$	$\lambda(\mu\text{m})$	$f_\nu(\text{Jy})$	$\lambda(\mu\text{m})$	$f_\nu(\text{Jy})$
1.6572E-02	1.2100E-07	1.6177E-02	1.8200E-09	1.6256E-02	4.7100E-08
1.8124E-02	1.7100E-07	1.7693E-02	2.7100E-09	1.7778E-02	6.6600E-08
1.9823E-02	2.4300E-07	1.9352E-02	4.0800E-09	1.9445E-02	9.4900E-08
2.1690E-02	3.3200E-07	2.1174E-02	5.8800E-09	2.1276E-02	1.3000E-07
2.3723E-02	4.4200E-07	2.3159E-02	8.2500E-09	2.3271E-02	1.7200E-07

NOTE. — The complete version of this table is in the electronic edition of the Journal. The printed edition contains only a sample.